

(19) World Intellectual Property
Organization
International Bureau



29 APR 2005 33 355



(43) International Publication Date
21 May 2004 (21.05.2004)

PCT

(10) International Publication Number
WO 2004/041778 A2

(51) International Patent Classification⁷:

C07D

(21) International Application Number:

PCT/US2003/034777

(22) International Filing Date: 31 October 2003 (31.10.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/422,986

1 November 2002 (01.11.2002) US

(71) Applicant (*for all designated States except US*): UNIVERSITY OF MEDICINE AND DENTISTRY OF NEW JERSEY [US/US]; 335 George Street, New Brunswick, NJ 08901 (US).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): BLACK, Ira, B. [US/US]; 158 Montgomery Road, Stillman, NJ 08558 (US).

(74) Agent: LICATA, Jane, Massey; Licata & Tyrrell P.C., 66 E. Main Street, Marlton, New Jersey 08053 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, ~~TJ~~, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A METHOD FOR INCREASING SYNAPTIC GROWTH OR PLASTICITY

(57) Abstract: The present invention relates to a method of modulating synaptic growth or plasticity by increasing the expression of genes found to be induced by BDNF. Such genes include c-fos proto-oncogene, early growth response protein 1, activity-regulated cytoskeletal associated, fos-related antigen 2, G1/S-specific cyclin D1, voltagegated potassium channel protein, sodium channel beta 1 subunit, secretogranin II precursor, somatostatin receptor 4, transmembrane receptor UNC5 homology, neuropeptide Y, VGF protein precursor and protein-tyrosine phosphatase 1B. Methods of identifying agents which modulate the expression of these genes and the use of such agents for treating a disease or condition associated with damaged or diseased synapses are also provided.

WO 2004/041778 A2

A METHOD FOR INCREASING SYNAPTIC GROWTH OR PLASTICITY

Background of the Invention

5 Neurotrophin-induced modifications of synaptic strength have been associated with mechanisms of learning and memory (Chao (2000) *J. Neurosci. Res.* 59:353-355; Tyler et al. (2002) *Learn. Mem.* 9:224-237). However, the molecular genetics of BDNF-induced plasticity is limited.

10 Several studies have drawn links between BDNF, learning, transcription and translation. For example, endogenous BDNF as well as transcriptional activation are necessary for the maintenance phase of long-term potentiation (L-LTP) (Korte et al. (1998) *Neuropharmacology* 37:553-559; Chen et al.

15 (1999) *J. Neurosci.* 19:7983-7990; Patterson et al. (2001) *Neuron* 32:123-140). In addition, BDNF and trkB may be required for the acquisition, consolidation and recall some types of information (Tyler et al. (2002) *supra*; Yamada et al. (2002) *Life Sci.* 70:735-744). Further, BDNF alone is

20 sufficient to trigger a form of long-term synaptic potentiation *in vivo* (BDNF-LTP) (Ying et al. (2002) *J. Neurosci.* 22:1532-1540), the induction of which requires transcription (Messaoudi et al., 2002). The regulation and function of specific candidate genes in BDNF-LTP, L-LTP and

25 learning paradigms have been demonstrated (Hevroni et al. (1998) *J. Mol. Neurosci.* 10:75-98; Guzowski et al. (2001) *J. Neurosci.* 21:5089-5098; Ying et al. (2002) *supra*). Although studies have employed transcriptional profiling to define panels of genes associated with learning (Luo et al.

30 (2001) *J. Mol. Neurosci.* 17:397-404; Cavallaro et al. (2002) *Neurochem. Res.* 27:1201-1207; Donahue et al. (2002) *Hippocampus* 12:821-833), the identification of a catalog of genes associated with BDNF-induced plasticity *in vitro*, and

-2-

the correlation of expression of these genes with learning in vivo has not been studied.

The regulated genes identified by transcriptional profiling serve as powerful tools to decipher mechanisms of both BDNF-induced transcription and plasticity. It has been shown that BDNF elicits differential electrophysiological responses in individual neurons (Levine et al. (1996) *Brain Res. Mol. Brain Res.* 38:300-303; Gottschalk et al. (1998) *J. Neurosci.* 18:6830-6839; Lessmann and Heumann (1998) *Neuroscience* 86:399-413; Schinder et al. (2000) *Neuron* 25:151-163; Thakker-Varia et al. (2001) *J. Neurosci.* 21:6782-6790). This phenomenon has been attributed to target specificity (Schinder et al. (2000) *supra*) and/or initial synaptic strength (Gottschalk et al. (1998) *supra*; Lessmann and Heumann (1998) *supra*; Berninger et al. (1999) *Learn Mem.* 6:232-242); however, these studies did not address the molecular mechanisms associated with distinctive single cell synaptic responses to BDNF. Moreover, while the signal transduction pathways required for BDNF-modulated synaptic plasticity have been examined (Gottschalk et al. (1999) *Learn. Mem.* 6:243-256; Minichiello et al. (2002) *Neuron* 36:121-137; Mizuno et al. (2003) *Mol. Psychiatry* 8:217-224), the requirement for various kinase pathways in BDNF-induced transcription remains to be determined.

Increased transcription during the processes of learning in vivo suggests that gene products may be involved in the formation of new memories. Transcriptional analysis following different learning tasks has revealed regulation of genes previously implicated in synaptic function (Guzowski et al. (2001) *supra*; Luo et al. (2001) *supra*; Cavallaro et al. (2002) *supra*; Donahue et al. (2002) *supra*).

-3-

There are only a few studies delineating functional roles for neuronal genes identified by transcriptional profiling (Huh et al. (2000) *Science* 290:2155-2159). Differential display has been used in conjunction with
5 electrophysiological analysis and gene-targeted mice to identify a requirement for Rab3A, a synaptic vesicle trafficking protein, in BDNF-induced plasticity (Thakker-Varia et al. (2001) *supra*).

10 Summary of the Invention

One aspect of the present invention is a method for modulating synaptic growth or plasticity. The method encompasses increasing the expression of a BDNF-inducible nucleic acid sequence, or activity of a protein encoded
15 thereby, so that synaptic growth or plasticity is stimulated. Preferably, the nucleic acid sequence which is inducible by BDNF is c-fos proto-oncogene (SEQ ID NO:1); early growth response protein 1 (SEQ ID NO:2); activity-regulated cytoskeletal associated (SEQ ID NO:3); fos-
20 related antigen 2 (SEQ ID NO:4); G1/S-specific cyclin D1 (SEQ ID NO:5); voltage-gated potassium channel protein (SEQ ID NO:6); sodium channel, beta 1 subunit (SEQ ID NO:7); secretogranin II precursor (SEQ ID NO:8); somatostatin receptor 4 (SEQ ID NO:9); transmembrane receptor UNC5
25 homology (SEQ ID NO:10); neuropeptide Y (SEQ ID NO:11); VGF protein precursor (SEQ ID NO:12); or protein-tyrosine phosphatase 1B (SEQ ID NO:13).

A nucleic acid sequence or protein of the invention is intended to include those disclosed herein (*i.e.*, SEQ ID
30 NO:1-13 and proteins encoded thereby) or any fragment, homolog, or ortholog which has the capacity to modulate synaptic growth or plasticity.

-4-

Another aspect of the present invention is a method for identifying an agent which increases synaptic growth or plasticity. The method involves contacting a test cell with an agent and detecting activation of c-fos proto-oncogene (SEQ ID NO:1); early growth response protein 1 (SEQ ID NO:2); activity-regulated cytoskeletal associated (SEQ ID NO:3); fos-related antigen 2 (SEQ ID NO:4); G1/S-specific cyclin D1 (SEQ ID NO:5); voltage-gated potassium channel protein (SEQ ID NO:6); sodium channel, beta 1 subunit (SEQ ID NO:7); secretogranin II precursor (SEQ ID NO:8); somatostatin receptor 4 (SEQ ID NO:9); transmembrane receptor UNC5 homology (SEQ ID NO:10); neuropeptide Y (SEQ ID NO:11); VGF protein precursor (SEQ ID NO:12); or protein-tyrosine phosphatase 1B (SEQ ID NO:13) nucleic acid sequences in the test cell. An increase in the activation of said nucleic acid sequences in the test cell contacted with the agent relative to the activation of said nucleic acid sequences in a test cell not contacted with the agent is indicative that said agent increases synaptic growth or plasticity.

A further aspect of the invention is a method for treating a disease or condition associated with damaged or diseased synapses. The method encompasses administering an effective amount of an agent which activates c-fos proto-oncogene (SEQ ID NO:1); early growth response protein 1 (SEQ ID NO:2); activity-regulated cytoskeletal associated (SEQ ID NO:3); fos-related antigen 2 (SEQ ID NO:4); G1/S-specific cyclin D1 (SEQ ID NO:5); voltage-gated potassium channel protein (SEQ ID NO:6); sodium channel, beta 1 subunit (SEQ ID NO:7); secretogranin II precursor (SEQ ID NO:8); somatostatin receptor 4 (SEQ ID NO:9); transmembrane receptor UNC5 homology (SEQ ID NO:10); neuropeptide Y (SEQ ID NO:11); VGF protein precursor (SEQ ID NO:12); or

-5-

protein-tyrosine phosphatase 1B (SEQ ID NO:13) nucleic acid sequences.

Brief Description of the Drawings

5 Figure 1 shows that Arc expression and synaptic response to BDNF correlate at the single-cell level. Figure 1A presents Arc expression in individual BDNF-treated cells following electrophysiological recordings and aRNA amplification. Arc transcription was measured by real-time
10 RT-PCR and then normalized to internal GAPDH levels. The data are then expressed as a ratio of the average Arc expression in the seven control cells (expression ratio). Figure 1B presents a correlation of Arc expression ratios to individual cell's synaptic BDNF responses. Average
15 synaptic charge 3 to 5 minutes post-BDNF exposure expressed as a fold increase over baseline (-2 to 0 minute) was plotted against Arc expression ratios shown in Figure 1A. Each circle represents one BDNF-treated cell. The cell number next to the circle corresponds with the cell number
20 in Figure 1A. Dashed line depicts a correlation coefficient indicating significance ($r = 0.66$, $p < 0.005$). The data are categorized into two groups of BDNF-treated cells: a cluster of non-responders (shaded) and a spread of responders (clear). Figure 1C shows the average Arc
25 expression ratios of the two populations of cells defined in Figure 1B. Responders (white bar) show a significantly higher fold increase in Arc expression (average \pm se, $n=7$) compared to non-responders (shaded bar) (average \pm se, $n=9$) (* $p<0.05$, t-test).

30 Figure 2 shows that VGF as well as immediate early genes are upregulated after training with trace eyeblink conditioning. RNA was obtained from the hippocampi of naïve rats, unpaired controls and trace conditioned rats

following 800 trials (200 trials/day for 4 days) of paired stimuli and subjected to real-time RT-PCR for a number of genes identified by microarray. Bars represent average gene expression \pm se for naïve rats ($n = 5$, white bar), unpaired
5 controls ($n=3$, stippled bar) and trace conditioning ($n=8$, black bar). All data is normalized to internal GAPDH levels and then expressed as a fold change over the amount of mRNA in naïve (control) rats. A number of immediate early genes including *c-fos*, *EGR1*, and *Arc* were upregulated after trace
10 conditioning. In addition, the secreted polypeptide VGF was increased following paired trace conditioning. *indicates paired samples are significantly different from unpaired and naïve control samples ($p < 0.05$, ANOVA).

15 Detailed Description of the Invention

Whole-cell patch-clamp recordings in conjunction with differential display and single-cell transcriptional analysis were used to identify known and novel genes with altered expression in BDNF-induced plasticity. The
20 functions of some of the identified genes have been defined using various methods.

The molecular genetics underlying BDNF-evoked plasticity were examined by expression profiling using cDNA microarrays on a population of hippocampal cells *in vitro*.
25 Synaptic effects of BDNF are observed within 20 minutes of exposure (Levine et al. (1996) *supra*; Lessmann and Heumann (1998) *supra*; Schinder et al. (2000) *supra*). A 3-hour BDNF treatment has been used to identify several genes with roles in synaptic plasticity (Thakker-Varia et al. (2001)
30 *supra*). Consequently, to reveal immediate, as well as downstream transcriptional changes, both acute (20 minutes) and longer treatment (3 hours) with BDNF were used herein. Approximately 8% of the arrayed cDNAs exhibited

-7-

differential expression at 20 minutes and 3 hours. Genes involved in synaptic plasticity and exhibiting an increase of 1.7-fold over control were selected for confirmation by real-time reverse transcription PCR (RT-PCR). Table 1 lists the genes induced by BDNF treatment. Two independent Atlas arrays were performed using RNA from hippocampal cells treated for either 20 minutes or 3 hours of BDNF (50 ng/mL) and control sister cultures.

TABLE 1

20 Minute BDNF Treatment					
Clone ID	Gene/Protein Name	Accession Number	Fold Change	NGF	SEQ ID NO:
A10g	c-fos	X06769	2.2 ± 0.4	UC	1
E06d	EGR1	M18416	4.3 ± 0.8	UC	2
F13I	Arc	U19866	13.7 ± 5.0	UC	3
3 Hour BDNF Treatment					
Clone ID	Gene/Protein Name	Accession Number	Fold Change	NGF	SEQ ID NO:
A05d	Fra2	U18913	2.3 ± 0.4	DOWN	4
A07a	CyclinD1	D14014	3.3 ± 0.5	UC	5
A10g	c-fos	X06769	2.3 ± 0.4	UC	1
B05d	Kv+1.1	M26161	1.4 ± 0.1	UC	6
B06j	Na+ch	M91808	1.8 ± 0.2	UC	7
B13n	SgII	M93669	5.1 ± 1.2	UC	8
D04i	SSTR4	U04738	1.7 ± 0.1	DOWN	9
D11k	UNC5H2	U87306	2.2 ± 0.6 [†]	UC	10
D13l	NPY	M20373	1.9 ± 0.3 [†]	DOWN	11
D14f	VGF	M60525	6.5 ± 1.5	UC	12
E06d	EGR1	M18416	6.6 ± 0.9	UC	2
E12f	PTP1B	M33962	1.4 ± 0.1	UC	13
F13l	Arc	U19866	34.8 ± 4.3	UC	3

10 The criteria for selection from the array was a >1.7-fold change. RT-PCR validated the induction of 16 genes (average ± standard error, n=6, [†]n=5, p<0.05, t-test). UC=unchanged, down=decreased.

15 Of those genes selected, the regulation of 7% at 20 minutes and 40% at 3 hours were validated, although the magnitude of regulation was not always identical to that observed on the microarray. To focus on genes specifically involved in plasticity, a related neurotrophin, nerve

-8-

growth factor (NGF), not inducing synaptic plasticity (Levine et al. (1996) *supra*), served as a control. NGF did not affect transcription of most genes, indicating the specificity of BDNF.

5 Genes upregulated by BDNF were categorized into different functional classes. At 20 minutes, immediate early genes were upregulated, whereas at 3 hours, in addition to those same immediate early genes, channel proteins, receptors and neuropeptides exhibited increases.

10 Two genes were of particular interest due to their association with synaptic plasticity: *Arc/Arg 3.1*, an effector of immediate early genes and *VGF*, a secreted polypeptide located in large dense core vesicles (Salton et al. (2000) *Front. Neuroendocrinol.* 21:199-219). *Arc*, the

15 most highly regulated gene at both 20 minutes and 3 hours, has been implicated in BDNF-induced plasticity and LTP (Guzowski et al. (2000) *J. Neurosci.* 20:3993-4001; Messaoudi et al. (2002) *J. Neurosci.* 22:7453-7461; Steward and Worley (2002) *Neurobiol. Learn. Mem.* 78:508-527; Ying

20 et al. (2002) *supra*). *VGF*, previously cloned from plate V of a NGF-induced PC12 cDNA library (Salton et al. (2000) *supra*), in contrast, was only increased at 3 hours and has not been associated with BDNF-induced synaptic efficacy. These two genes were used in subsequent experiments as

25 indices of synaptic efficacy and prototypes for functional studies.

To examine the relationship between synaptic response and gene expression, an individual cell's electrophysiological response to BDNF was examined in

30 conjunction with single-cell RT-PCR for *Arc*. Previous studies revealed a range of synaptic activities within a population of pyramidal-like hippocampal neurons following 3 hours BDNF exposure (Thakker-Varia et al. (2001) *supra*).

Whole-cell patch-clamp recordings from 16 cells verified 0.8- to 4.5-fold increase in synaptic charge following 20 minutes BDNF exposure. Average synaptic charge for control cells not exposed to BDNF (n=7) exhibited no change in synaptic charge during the length of the recording. A group of BDNF-treated cells presenting a minimal synaptic response was apparent. This group was categorized as non-responders (Figure 1B) and the other cells showing a greater response to BDNF were classified as responders. To determine whether cellular basal activity influences the response to BDNF, the average synaptic charge pre-BDNF of the responders (36.3 ± 13.1 pC, n=7) versus the non-responders (23.2 ± 6.6 pC, n=9) was compared. There was no statistical difference in the populations ($p > 0.05$, t-test) examined between initial synaptic strength and responsiveness to BDNF, indicating that this parameter is not responsible for the differential response.

The molecular genetics of differential synaptic responses to BDNF was examined by performing single-cell transcriptional analysis after whole-cell patch-clamp. Arc was selected as a prototype due to its acute and robust induction by BDNF in a population of hippocampal neurons. To assess the relationship between differential electrophysiological responses to BDNF and Arc gene expression, individual cells were harvested following electrophysiological recordings. Figure 1A represents Arc expression in cells exposed to BDNF relative to the average Arc expression in cells not exposed to BDNF (control) as assessed by real-time RT-PCR (expression ratio). A range of Arc expression levels was apparent within the group of 16 BDNF-treated cells, indicating transcriptional differences among pyramidal-like hippocampal cells.

-10-

The relationship between the synaptic response to BDNF and transcriptional changes was examined. Synaptic charge was plotted against Arc expression ratios for individual cells (Figure 1B). A linear correlation ($r=0.66$, $p<0.005$) was evident as depicted by the dashed line, indicating a direct relationship between synaptic plasticity and gene expression. Alternatively, the data may be viewed as representing two sets of BDNF-treated cells: a cluster of non-responders (shaded area) and a spread of responders (clear area). Classification of cells into electrophysiologic responders or non-responders, revealed a striking difference in Arc expression ratios (Figure 1C). Responders exhibit a 17-fold increase in Arc expression relative to the non-responders ($p<0.05$, t -test). These results demonstrate that the magnitude of BDNF-induced synaptic plasticity is reflected in gene expression in individual hippocampal cells. Enhanced transcription following neurotrophin exposure, may represent a mechanism for long-term maintenance of increased synaptic activity as has been shown for L-LTP (Nguyen et al. (1994) *Science* 265:1104-1107).

Signaling cascades involved in BDNF-mediated synaptic strengthening include mitogen-activated protein (MAP) kinase, phospholipase C- γ (PLC- γ) and phosphoinositide-3 kinase (PI3K) (Gottschalk et al. (1999) *supra*; Minichiello et al. (2002) *supra*; Ying et al. (2002); Mizuno et al. (2003) *supra*). To determine whether these transduction pathways are also required for BDNF transcriptional regulation of immediate early genes at 20 minutes and for *VGF* at 3 hours, pharmacological inhibitors were employed (Table 2).

TABLE 2

Inhibitor (Protein Inhibited)	20 Minutes			3 Hours
	c-fos	EGR-1	Arc	VGF
K252a (Tyr Kinase)	Blocked	Blocked	Blocked	Blocked
U0126 (MEK)	Blocked	Blocked	Blocked	Blocked
PD98059 (MEK)	Blocked	Blocked	Blocked	Blocked
U73122 (PLC- γ)	Blocked	UC	Blocked	Blocked
LY294002 (PI3 Kinase)	UC	UC	UC	UC
KN93 (CaMKII)	Blocked	Blocked	Blocked	Blocked
H-89 (PKA)	UC	UC	UC	UC
CC (PKC)	UC	UC	UC	UC

Effect of inhibitors on BDNF-induced gene transcription. The regulation of BDNF-induced genes in the presence of inhibitors was determined by real time RT-PCR. Neurons were pre-incubated with inhibitors for 30 minutes prior to BDNF treatment (20 minutes for immediate early genes and 3 hours for VGF). One group of cells in each experiment was treated with inhibitor alone and showed no significant change in expression relative to vehicle control. "blocked" indicates significant reduction of transcription in the presence of inhibitor plus BDNF relative to BDNF alone ($p < 0.05$, ANOVA, $n=3$). UC (unchanged) indicates no significant effect of inhibitor on transcription.

Each experiment consisted of four groups: vehicle, inhibitor alone, BDNF, BDNF plus inhibitor. Data were normalized to GAPDH and expressed as a fold change of vehicle control. Inhibitor alone showed no significant effect on transcription. K252a, U0126, PD98059, U73122 and KN93 blocked BDNF-induced transcription of the immediate early genes and VGF whereas LY294002, H-89 and chelerythrine chloride (CC) did not affect gene expression. K252a blocked the BDNF effect, indicating that transcription results from trkB stimulation. Pretreatment with either MAP kinase inhibitor (U0126 or PD98059) blocked BDNF-induced transcription, indicating a requirement for

-12-

MEK-mediated phosphorylation of MAP kinase. The PLC- γ inhibitor (U73122) and the Ca²⁺/calmodulin-dependent protein kinase (CaMK) inhibitor (KN93) blocked transcription of most immediate early genes and VGF (Table 2). In contrast, a number of signaling pathways were not involved in BDNF-induced transcription, shown by the lack of effect of inhibitors of PI3K (LY294002), protein kinase A (PKA) (H-89) and protein kinase C (PKC) (chelerythrine chloride, CC). These results indicate that BDNF-mediated transcription at both 20 minutes and 3 hours was dependent on the CaMK and PLC- γ pathways in addition to trkB and MAP kinase.

To determine whether VGF and other transcripts induced by BDNF *in vitro* predict alterations in mRNA *in vivo*, gene expression was examined following training on a hippocampal-dependent learning task. Rats were exposed to 800 trials of either paired or unpaired trace eyeblink conditioning and sacrificed 24 hours later. Eyeblinks were assessed by eyelid electromyography and those occurring during the 500 ms trace interval were considered conditioned responses (CRs). Rats exposed to paired stimuli exhibited 60% (± 4) CRs, whereas those exposed to the same number of stimuli presented in an explicitly unpaired manner exhibited 11% (± 1) CRs, a percentage similar to that of spontaneous blinks (10%). A t-test for independent samples on the number of conditioned responses revealed that the difference was significant at $p < 0.0001$. VGF transcripts were significantly upregulated in the hippocampus after paired but not unpaired eyeblink conditioning or naïve controls ($p < 0.05$, ANOVA) (Figure 2). Several of the immediate early genes including *c-fos*, *EGR1* and *Arc* were also induced by exposure to the paired stimuli

-13-

($p < 0.05$, ANOVA). These changes were specific and selective, since other genes induced by BDNF *in vitro* were not upregulated by trace eyeblink conditioning; *Na⁺ channel*, *somatostatin receptor 4 (SSTR4)* and *transmembrane receptor UNC homolog 2 (UNC5H2)* were not significantly enhanced following paired eyeblink training. These results indicate that BDNF-induced transcriptional alterations of *VGF* and immediate early genes *in vitro* predict changes *in vivo* in the hippocampus during memory formation. Accordingly, modulating the neuropeptide *VGF*, for example, is useful for regulating synaptic function associated with learning and memory.

The products of the genes induced following learning may regulate neuronal plasticity at the single-cell level. *VGF*, like BDNF, is a secreted polypeptide and thus may be relevant for intercellular communication. A functional role for *VGF* in synaptic plasticity was thus investigated. To determine whether BDNF-induced transcriptional changes of *VGF* were accompanied by translational alterations, *VGF* levels were examined following BDNF treatment by western blot analysis. *VGF*, migrating as a doublet with an apparent molecular weight of 85-90 kD, increased by 1.5-fold within 3 hours of BDNF exposure. *VGF* protein levels were maximal at 12 hours and remained elevated for 48 hours of continuous BDNF application. Consequently, increased *VGF* mRNA levels after 3 hours BDNF treatment were associated with enhanced protein translation. *VGF* protein processing occurs in both neurons and neuroendocrine cells (Trani et al. (1995) *J. Neurochem.* 65:2441- 2449; Salton et al. (2000) *supra*). The localization of *VGF* in dissociated hippocampal cultures was examined using a C-terminal antibody (R-15) which labeled both the soma and neurites indicating translocation to the neuronal processes. The

-14-

expression of VGF was similar to that previously observed in primary neuronal cultures (Benson and Salton (1996) *Brain Res. Dev. Brain Res.* 96:219-228) and was punctate in appearance. This pattern is consistent with the regulated
5 release of VGF from large dense core vesicles (Trani et al. (1995) *supra*; Benson and Salton (1996) *supra*; Possenti et al. (1999) *Endocrinology* 140:3727-3735) and indicates a physiological role for the peptide.

To explore a functional role for VGF in
10 neuromodulation, whole-cell patch-clamp recordings on hippocampal cells were performed in the presence of synthetic VGF peptides. The electrophysiological profile of VGF peptides on hippocampal cells was similar to that observed for BDNF (Levine et al. (1996) *supra*). Increased
15 synaptic charge was observed within 5 minutes of perfusion of peptides derived from the C-terminal portion of VGF (Thr-Leu-Gln-Pro-62; SEQ ID NO:14), whereas no change in synaptic charge resulted from application of peptide from the N-terminal region (Leu-Glu-Gly-Ser-28; SEQ ID NO:15).
20 Enhanced synaptic activity was maintained for up to 25 minutes in the continued presence of Thr-Leu-Gln-Pro-62 (SEQ ID NO:14). As an independent confirmation of the effect of VGF on synaptic plasticity, an alternate peptide derived from the C-terminal region was tested (Ala-Gln-Glu-Glu-30; SEQ ID NO:16) and revealed a similar effect on
25 synaptic charge. The effects of both Ala-Gln-Glu-Glu-30 (SEQ ID NO:16) and Thr-Leu-Gln-Pro-62 (SEQ ID NO:14) peptides were dose-dependent, saturating at 0.1 μ M. Taken together, these data indicate that the functional portion
30 of VGF resides in the C-terminal region of the protein. These findings are the first demonstration of a neuromodulatory role for VGF in synaptic plasticity of the hippocampus which subserves learning and memory.

-15-

The present invention relates to differentially-expressed genes involved in BDNF regulation of synaptic activity. These genes, referred to herein as nucleic acid sequences of the invention, which were inducible by BDNF

5 include c-fos proto-oncogene (c-fos; SEQ ID NO:1); early growth response protein 1 (EGR1; SEQ ID NO:2); activity-regulated cytoskeletal associated (Arc; SEQ ID NO:3); fos-related antigen 2 (Fra2; SEQ ID NO:4); G1/S-specific cyclin D1 (cyclinD1; SEQ ID NO:5); voltage-gated potassium channel

10 protein (Kv+ 1.1; SEQ ID NO:6); sodium channel, beta 1 subunit (Na+ ch; SEQ ID NO:7); secretogranin II precursor (SgII; SEQ ID NO:8); somatostatin receptor 4 (SSTR4; SEQ ID NO:9); transmembrane receptor UNC5 homology (UNC5H2; SEQ ID NO:10); neuropeptide Y (NPY; SEQ ID NO:11); VGF protein

15 precursor (VGF; SEQ ID NO:12); and protein-tyrosine phosphatase 1B (PTP1B; SEQ ID NO:13). Said genes were found to be upregulated or activated in BDNF-induced plasticity. Accordingly, as these genes may contribute to synaptic growth and plasticity, agents which modulate the expression

20 or activity of these genes are useful for modulating learning and memory. As used herein, synaptic growth and plasticity is intended to include both acute and long-term activation, presynaptic and postsynaptic, and basal and enhanced transmission.

25 Accordingly, one aspect of the present invention is a method for increasing synaptic growth or plasticity via activating or increasing the expression of a nucleic acid sequence, which is inducible by BDNF, or the activity of a protein encoded thereby. Such an increase in gene

30 expression or protein activity is useful in promoting or stimulating synaptic growth or plasticity to promote or stimulate, for example, memory or learning. Means of increasing the expression of a nucleic acid sequence

-16-

disclosed herein or the activity of proteins encoded thereby are provided.

It is contemplated that the expression of a BDNF-induced nucleic acid sequence of the invention may be increased via genetic engineering (gene therapy) or by administering agents which increase the expression or activity of said nucleic acid sequences or proteins encoded thereby, respectively. It is further contemplated that a protein of the invention may be administered as a purified recombinant protein (e.g., encapsulated in liposome formulations) to increase the amount of said protein in a cell. In particular it is advantageous for the gene, agent or protein to be expressed or administered to cells of the brain, e.g., the hippocampus.

In one embodiment, genetic engineering or gene therapy approaches to increase the expression of a nucleic acid sequence of the invention to promote learning or memory may employ any method known in the art. For example, a method for increasing expression of a nucleic acid sequence of the invention includes exogenously supplying said nucleic acid sequence to a cell (i.e., a gene knockin).

In general, the coding sequence of a nucleic acid sequence of the invention (i.e., mRNA or DNA sequence) is delivered to cells in an area in which increased memory or learning is desirable. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, may be used. Alternatively, if it is desired that the cells stably retain the construct, it may be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. Additionally, the construct may be introduced into a neuronal cell (e.g., cortical neurons or dorsal root ganglion neurons) or a cell

-17-

capable of differentiating into a neuronal cell (e.g., stem cell) and be transplanted into an individual in need. The construct may include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and
5 a transcriptional terminator signal, for controlling transcription of the coding sequence in the cells.

A DNA construct or RNA molecule encoding a nucleic acid sequence of the invention may be directly introduced into a cell using well-known methods or incorporated into a
10 viral vector for delivery to the host animal. Vectors, such as replication-defective retroviruses, rabies virus, adenoviruses and adeno-associated viruses may be used, preferably a neurotropic virus is used. Protocols for producing recombinant viruses and for infecting cells *in*
15 *vitro* or *in vivo* with such viruses are described by in Ruitenberg, et al. ((2002) *Methods* 28(2):182-94); Lundstrom ((2001) *Curr. Gene Ther.* 1(1):19-29) and Current Protocols in Molecular Biology, Ausubel, F. M. et al. (eds.) Greene Publishing Associates, (1989), Sections 9.10-9.14 and other
20 standard laboratory manuals. The genome of an adenovirus may be manipulated such that it encodes and expresses a nucleic acid sequence of the invention but is inactivated in terms of its ability to replicate in a normal lytic viral life cycle (Berkner, et al. (1988) *BioTechniques*
25 6:616; Rosenfeld, et al. (1991) *Science* 252:431-434; Rosenfeld, et al. (1992) *Cell* 68:143-155). Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 dl324 or other strains of adenovirus (e.g., Ad2, Ad3, Ad7, etc.) are well-known to those skilled in the art.
30 *In vivo* use of adenoviral vectors is described in Flotte, et al. ((1993) *Proc. Natl. Acad. Sci. USA* 90:10613-10617) and Kaplitt, et al. ((1994) *Nature Genet.* 8:148-153). Other viral vectors, such as those based on togaviruses, alpha

-18-

viruses, or vaccinia virus may also be used. Alternatively, an adeno-associated virus vector such as that disclosed by Xu, et al. ((2001) *Gene Ther.* 8(17):1323-32) may be used to express a nucleic acid sequence of the invention.

5 Regulatory sequences for *in vivo* expression of a nucleic acid sequence of the invention are preferably brain-specific and may be constitutive or regulated. Exemplary regulatory sequences include those associated with the expression of prions, human glial fibrillary
10 acidic protein gene (Shi, et al. (2001) *Proc. Natl. Acad. Sci. USA* 98(22):12754-9), neuron-specific enolase or beta-actin.

A DNA construct or RNA molecule of the invention may also be combined with a condensing agent, such as
15 polylysine, polyarginine, polyornithine, protamine, spermine, spermidine, or putrescine, to form a gene delivery vehicle. Many suitable methods for making such linkages are known in the art. Alternatively, a DNA construct or RNA molecule may be associated with a liposome
20 for delivery to a desired cell. For example, plasmid DNA may be targeted to the brain with pegylated immunoliposomes using a targeting ligand such as a peptidomimetic mAb, which binds to a transporting receptor on the blood-brain barrier (Shi, et al. (2001) *Proc. Natl. Acad. Sci. USA*
25 98(22):12754-9). Other suitable methods of providing DNA constructs or RNA molecules include DNA-ligand combinations. Receptor-mediated DNA delivery techniques are also well-known in the art (Findeis, et al. (1993) *Trends Biotech.* 11:202-05; Chiou, et al. (1994) *Gene Therapeutics: Methods and Applications of Direct Gene Transfer* (J. A. Wolff, ed.); Wu, et al. (1994) *J. Biol. Chem.* 269:542-46).
30 Expression of a nucleic acid sequence of the invention may be monitored by detecting production of mRNA which

-19-

hybridizes to a delivered coding sequence or by detecting the protein product of the gene using, for example, immunological techniques.

Expression of an endogenous nucleic acid sequence of the invention in a cell may also be altered by introducing in-frame with the endogenous gene a DNA construct comprising a targeting sequence, a regulatory sequence, an exon, and an unpaired splice donor site by homologous recombination, such that a homologously recombinant cell comprising a new transcription unit is formed. The new transcription unit may be used to turn expression of a nucleic acid sequence of the invention on or off as desired. This method of affecting endogenous gene expression is taught in U.S. Patent No. 5,641,670.

Alternatively, expression of a nucleic acid sequence of the invention may be altered in cells which have been removed from a mammal, such as neuronal cells. The cells may then be replaced into the same or another mammal, preferably to or within the vicinity of a region which learning or memory is to be increased.

In another embodiment, a recombinantly-produced or chemically-synthesized protein of the invention may be used to increase said protein levels in a cell where it is desirable to increase or stimulate learning or memory. Proteins of the invention may be used in a pharmaceutically acceptable composition and may be administered intraperitoneally, intracranially, subcutaneously, in accordance with well-known protocols.

As will be appreciated by one of skill in the art, a full-length protein of the invention may be produced for use in the methods of the invention, however, fragments of a protein of the invention may also be used provided the fragment maintains the desired binding interaction or

-20-

activity of the full-length protein. Exemplary fragments of VGF which may be used to modulate synaptic growth or plasticity include fragments from the C-terminal region of VGF, e.g., Ala-Gln-Glu-Glu-30 (SEQ ID NO:16) and Thr-Leu-
5 Gln-Pro-62 (SEQ ID NO:14) peptides.

In general, recombinant production of a protein of the invention requires incorporation of nucleic acid sequences encoding said protein (e.g., SEQ ID NO:1-13) into a recombinant expression vector in a form suitable for
10 expression of the protein in a host cell.

A suitable form for expression provides that the recombinant expression vector includes one or more regulatory sequences operatively-linked to a nucleic acid sequence encoding a protein of the invention in a manner
15 which allows for transcription of the nucleic acids into mRNA and translation of the mRNA into the protein. Regulatory sequences may include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are known to those
20 skilled in the art and are described in Goeddel D.D., ed., Gene Expression Technology, Academic Press, San Diego, CA (1991). It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transfected and/or the level of
25 expression required. Nucleic acid sequences or expression vectors harboring nucleic acid sequences encoding a protein of the invention may be introduced into a host cell, which may be of eukaryotic or prokaryotic origin, by standard techniques for transforming cells. Suitable methods for
30 transforming host cells may be found in Sambrook, et al. (Molecular Cloning: A Laboratory Manual, 3rd Edition, Cold Spring Harbor Laboratory Press (2000)) and other laboratory manuals. The number of host cells transformed with a

-21-

nucleic acid sequence encoding a protein of the invention will depend, at least in part, upon the type of recombinant expression vector used and the type of transformation technique used. Nucleic acids may be introduced into a host cell transiently, or more typically, for long-term expression of a protein of the invention the nucleic acid sequence is stably integrated into the genome of the host cell or remains as a stable episome in the host cell. Once produced, a protein of the invention may be recovered from culture medium as a secreted polypeptide, although it also may be recovered from host cell lysates when directly expressed without a secretory signal. When a protein of the invention is expressed in a recombinant cell other than one of human origin, said protein is substantially free of proteins or polypeptides of human origin. However, it may be necessary to purify the protein from recombinant cell proteins or polypeptides using conventional protein purification methods to obtain preparations that are substantially homogeneous as to a protein of the invention. As a first step, the culture medium or lysate is centrifuged to remove particulate cell debris. The membrane and soluble protein fractions are then separated. The recombinant protein may then be purified from the soluble protein fraction. The recombinant protein thereafter is purified from contaminant soluble proteins and polypeptides using any of the following suitable purification procedures: by fractionation on immunoaffinity or ion-exchange columns; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, SEPHADEX G-75; ligand affinity chromatography, and protein A SEPHAROSE columns to remove contaminants such as IgG.

-22-

In addition to recombinant production, a protein of the invention may be produced by direct peptide synthesis using solid-phase techniques (Merrifield J. (1963) *J. Am. Chem. Soc.* 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer, Boston, MA). Various fragments of a protein of the invention may be chemically-synthesized separately and combined using chemical methods to produce a full-length molecule.

In a further embodiment of the present invention, expression of a nucleic acid sequence of the invention or activity of a protein encoded thereby may be regulated using an agent which increases the expression or activity of a protein of the invention. Agents suitable for regulating the expression or activity of protein of the invention may be identified in accordance with the screening assay of the invention and encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Agents may also be found among biomolecules including peptides, antibodies, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives,

-23-

or proteins, DNA constructs or RNA molecules described herein and structural analogs or combinations thereof.

Agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. The
5 use of replicable genetic packages, such as the bacteriophages, is one method of generating novel polypeptide entities that regulate the expression or activity of a protein of the invention. This method generally consists of introducing a novel, exogenous DNA
10 segment into the genome of a bacteriophage (or other amplifiable genetic package) so that the polypeptide encoded by the non-native DNA appears on the surface of the phage. When the inserted DNA contains sequence diversity, then each recipient phage displays one variant of the
15 template amino acid sequence encoded by the DNA, and the phage population (library) displays a vast number of different but related amino acid sequences.

Antibodies which specifically bind a protein of the invention are also contemplated as agonistic agents for
20 regulating the activity of a protein of the invention. Antibodies to a protein of the invention may be generated using methods that are well-known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments, and
25 fragments produced by a Fab expression library. Neutralizing antibodies are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others,
30 may be immunized by injection with a protein of the invention or any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological

-24-

response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface-active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole
5 limpet hemocyanin, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum* are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to a protein of the
10 invention have an amino acid sequence consisting of at least five amino acids and more preferably at least 10 amino acids. It is also preferable that they are identical to a portion of the amino acid sequence of the natural protein, and they may contain the entire amino acid
15 sequence of a small, naturally occurring molecule. Short stretches of amino acids of a protein of the invention may be fused with those of another protein such as keyhole limpet hemocyanin and antibody produced against the chimeric molecule.

20 Monoclonal antibodies to a protein of the invention may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique,
25 and the EBV-hybridoma technique (Kohler, et al. (1975) *Nature* 256:495-497; Kozbor, et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, et al. (1983) *Proc. Natl. Acad. Sci.* 80:2026-2030; Cole, et al. (1984) *Mol. Cell Biol.* 62:109-120).

30 In addition, techniques developed for the production of humanized and chimeric antibodies, the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological

-25-

activity may be used (Morrison, et al. (1984) *Proc. Natl. Acad. Sci.* 81, 6851-6855; Neuberger, et al. (1984) *Nature* 312:604-608; Takeda, et al. (1985) *Nature* 314:452-454). Alternatively, techniques described for the production of
5 single chain antibodies may be adapted, using methods known in the art, to produce specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton
10 (1991) *Proc. Natl. Acad. Sci.* 88,11120-11123).

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as is well-known in the art (Orlandi, et
15 al. (1989) *Proc. Natl. Acad. Sci.* 86: 3833-3837; Winter, et al. (1991) *Nature* 349:293-299).

Antibody fragments, which contain specific binding sites for a protein of the invention, may also be generated. For example, such fragments include, but are not
20 limited to, the $F(ab')_2$ fragments which may be produced by pepsin digestion of the antibody molecule and the Fab fragments which may be generated by reducing the disulfide bridges of the $F(ab')_2$ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and
25 easy identification of monoclonal Fab fragments with the desired specificity (Huse, et al. (1989) *Science* 254:1275-1281).

Various immunoassays may be used for screening to identify antibodies having the desired specificity.
30 Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificity are well-known in the art. Such immunoassays typically involve

the measurement of complex formation between a specific antibody and a protein of the invention. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes is preferred, but a competitive binding assay may also be employed.

While isolated proteins, DNA constructs, RNA molecules, or antibodies disclosed heretofore may be used for increasing the expression or activity of a protein of the invention so that synaptic growth or plasticity is modulated, the present invention also provides a method for identifying other agents for such purposes. A cell-based assay is specifically contemplated. Such a method involves contacting a test cell with an agent and detecting expression or activation of a nucleic acid sequence of the invention in the test cell. An agent which causes an increase in activation of a nucleic acid sequence of the invention in the test cell when compared to a test cell not contacted with the agent, indicates that the agent increases synaptic growth or plasticity.

One embodiment of the screening method of the invention is based on the use of a reporter to detect gene expression. In this embodiment, test cells are typically isolated neurons which contain or have been transformed with a nucleic acid sequence encoding a reporter operably linked to promoter of a nucleic acid sequence of the invention. Upon exposure to the test agent, the presence or activity of the reporter is detected and correlated with the expression or activation of a nucleic acid sequence of the invention.

A nucleic acid sequence encoding a reporter may be inserted into a recombinant expression vector to generate a reporter gene construct. A reporter gene construct refers

-27-

to a plasmid, virus or other vehicle known in the art that has been manipulated by insertion or incorporation of nucleic acid sequences encoding a reporter. Such reporter gene constructs of the invention are preferably plasmids which contain at least a portion of a promoter sequence of a nucleic acid sequence of the invention which is operably associated with the inserted nucleic acid sequences encoding the reporter. The construct typically contains an origin of replication as well as specific selectable or screenable marker genes for initially isolating, identifying or tracking test cells that contain the desired reporter/promoter DNA. The reporter gene construct also may provide unique or conveniently located restriction sites to allow severing and/or rearranging portions of the DNA inserts in a reporter gene construct. More than one reporter gene may be inserted into the construct such that the test cells containing the resulting construct may be assayed by different means.

A reporter refers to any sequence that is detectable and distinguishable from other sequences present in test cells. Preferably, the reporter nucleic acid sequence encodes a protein that is readily detectable either by its presence, or by its activity that results in the generation of a detectable signal. A nucleic acid sequence encoding the reporter is used in the invention to monitor and report the activity of a promoter of a nucleic acid sequence of the invention in test cells.

A variety of enzymes may be used as reporters including, but are not limited to, β -galactosidase (Nolan, et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:2603-2607), chloramphenicol acetyltransferase (CAT; Gorman, et al. (1982) *Mol. Cell Biol.* 2:1044; Prost, et al. (1986) *Gene*

45:107-111), β -lactamase, β -glucuronidase and alkaline phosphatase (Berger, et al. (1988) *Gene* 66:1-10; Cullen, et al. (1992) *Meth. Enzymol.* 216:362-368). Transcription of the nucleic acid sequences encoding a reporter leads to
5 production of the enzyme in test cells. The amount of enzyme present may be measured via its enzymatic action on a substrate resulting in the formation of a detectable reaction product. The method of the invention provides means for determining the amount of reaction product,
10 wherein the amount of reaction product generated or the remaining amount of substrate is related to the amount of enzyme activity. For some enzymes, such as β -galactosidase, β -glucuronidase and β -lactamase, well-known fluorogenic substrates are available that allow the enzyme to convert
15 such substrates into detectable fluorescent products.

A variety of bioluminescent, chemiluminescent and fluorescent proteins also may be used as light-emitting reporters. Exemplary light-emitting reporters, which are enzymes and require cofactor(s) to emit light, include, but
20 are not limited to, the bacterial luciferase (*luxAB* gene product) of *Vibrio harveyi* (Karp (1989) *Biochim. Biophys. Acta* 1007:84-90; Stewart, et al. (1992) *J. Gen. Microbiol.* 138:1289-1300), and the luciferase from firefly, *Photinus pyralis* (De Wet, et al. (1987) *Mol. Cell. Biol.* 7:725-737).

25 Another type of light-emitting reporter, which does not require substrates or cofactors includes, but is not limited to, the wild-type green fluorescent protein (GFP) of *Victoria aequoria* (Chalfie, et al. (1994) *Science* 263:802-805), modified GFPs (Heim, et al. (1995) *Nature* 373:663-4; WO 96/23810), and the gene products encoded by
30 the *Photobacterium luminescens lux* operon (*luxABCDE*) (Francis, et al. (2000) *Infect. Immun.* 68(6):3594-600).

-29-

Transcription and translation of these type of reporters leads to the accumulation of the fluorescent or bioluminescent proteins in test cells, which may be measured by a device, such as a fluorimeter, flow
5 cytometer, or luminometer. Methods for performing assays on fluorescent materials are well-known in the art (e.g., Lackowicz (1983) In: Principles of Fluorescence Spectroscopy, New York, Plenum Press).

For convenience and efficiency, enzymatic reporters
10 and light-emitting reporters are preferred for the screening assays of the invention. Accordingly, the invention encompasses histochemical, colorimetric and fluorometric assays.

A promoter which is operably associated or operably
15 linked to nucleic acid sequences encoding a reporter means that the sequences are joined and positioned in such a way as to permit transcription. Two or more sequences, such as a promoter and any other nucleic acid sequences are operably associated if transcription commencing in the
20 promoter will produce an RNA transcript of the operably associated sequences. A promoter is defined as a fully functional sequence which regulates expression of a nucleic acid sequence of the invention, i.e., a promoter which has binding sites for transcription factors and minimal
25 promoter sequences. A promoter may include sequences upstream of a coding region of SEQ ID NO:1-13.

Various portions of a promoter or gene sequence may be generated by PCR (e.g., by inverse PCR methods using primers based on the 5' sequences of SEQ ID NO:1-13 and
30 genomic DNA as a template) or other conventional cloning techniques using DNA from the genomic locus of a gene identified herein. For PCR amplification, primers may be synthesized corresponding to the 5' and 3' boundaries of

- 30 -

the selected promoter or gene regions. Primers also may contain additional restriction enzyme recognition sequences to facilitate subcloning.

5 Introduction of the reporter gene construct into the test cells may be carried out by conventional techniques well-known to those skilled in the art, such as transfection, transformation, conjugation, and transduction.

10 In addition to conventional chemical methods of transformation, the reporter gene construct of the invention may be introduced into a test cell by physical means, such as by electroporation or microinjection. Electroporation allows transfer of the vector by high voltage electric impulse, which creates pores in the plasma
15 membrane of the cell and is performed according to methods well-known in the art. Additionally, the reporter gene construct may be introduced into test cells by protoplast fusion, using methods well-known in the art. The reporter gene construct may be introduced into a test cell
20 transiently, or more typically, the nucleic acids are stably integrated into the genome of the test cell or remain as stable episomes in the test cell.

The test cells which contain the nucleic acid sequences encoding the reporter and which express products
25 of the nucleic acid sequences encoding the reporter may be identified by at least four general approaches; detecting DNA-DNA or DNA-RNA hybridization; observing the presence or absence of marker gene functions (e.g., resistance to antibiotics); assessing the level of transcription as
30 measured by the expression of reporter mRNA transcripts in the host cell; and detecting the reporter gene product as measured by immunoassay or by its biological activity.

-31-

The steps involved in a reporter-based screening assay include, culturing a test cell which contains nucleic acid sequences encoding a reporter operably linked to a promoter of a nucleic acid sequence of the invention; adding at
5 least one test agent to a point of application, such as a well, in the plate containing the test cell and incubating the plate for a time sufficient to allow the test agent to effect reporter accumulation; detecting reporter activity of the test cell contacted with the test agent, wherein
10 reporter activity indicates expression of the reporter polypeptide in the test cell; and comparing reporter activity of the test cell which has been contacted with the test agent to that of the test cell not contacted with the test agent. An increase in reporter activity of the test
15 cell contacting the test agent relative to the reporter activity of the test cell not contacting the test agent indicates that the test agent causes an increase in expression of a nucleic acid sequence of the invention in the test cell.

20 Another embodiment of the screening method of the invention is based on the detection of gene expression using RT-PCR and/or microarray analysis. In this embodiment, at least one test agent is added to a test cell located, for example, in a well of a microtiter plate, and
25 the test cell and test agent are incubated for a time sufficient to allow the test agent to effect accumulation of a nucleic acid sequence of the invention. Subsequently, expression or activation of said nucleic acid sequence is determined by reverse-transcription (RT) alone or in
30 combination with array hybridization as disclosed herein. An increase in expression of said nucleic acid sequence in the test cell contacting the test agent, relative to expression of the same in a test cell not contacting the

-32-

test agent, indicates that the test agent is useful for modulating synaptic growth or plasticity.

Test cells used in accordance with a screening method of the invention may be cultured under standard conditions of temperature, incubation time, optical density, plating density and media composition corresponding to the nutritional and physiological requirements of the cells. However, conditions for maintenance and growth of the test cell may be different from those for assaying candidate agents in the screening methods of the invention. Modified culture conditions and media are used to facilitate detection of the expression of a reporter molecule. Any techniques known in the art may be applied to establish the optimal conditions.

Screening assays of the invention may be performed in any format that allows rapid preparation and processing of multiple reactions such as in, for example, multi-well plates of the 96-well variety. Stock solutions of the agents as well as assay components are prepared manually and all subsequent pipeting, diluting, mixing, washing, incubating, sample readout and data collecting is done using commercially available robotic pipeting equipment, automated work stations, and analytical instruments for detecting the signal generated by the assay.

In addition to the agent and a test cell, a variety of other reagents may be included in the screening assays. These include reagents like salts, neutral proteins, e.g., albumin, detergents, etc. Also, reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, and the like may be used.

Activators or repressors of expression of a nucleic acid sequence of the invention may include agents which

-33-

directly bind to a region on the promoter as well as agents which interact with transcriptional factors that bind to the said promoter thereby specifically regulating gene expression.

5 Agents identified in the screening assays provided herein are useful in stimulating or increasing synaptic growth or plasticity. Thus, these agents, as well as isolated proteins of the invention, or fragments thereof, may be formulated into pharmaceutical compositions
10 comprising an effective amount of a recombinant protein or agent to increase the expression, protein levels, or activity of the inventive nucleic acid sequences or proteins encoded thereby. Such pharmaceutical compositions may be prepared by methods and contain vehicles which are
15 well-known in the art. A generally recognized compendium of such methods and ingredients is Remington: The Science and Practice of Pharmacy, Alfonso R. Gennaro, editor, 20th ed. Lippincott Williams & Wilkins: Philadelphia, PA, 2000.

By effective amount it is meant an amount of
20 recombinant protein or agent which stimulates or increases synaptic growth or plasticity by 25%, 50%, 75%, or more compared to individuals not receiving treatment. Preferably, the agent or protein provides both therapeutic treatment and prophylactic or preventative measures,
25 wherein the object is to prevent or slow down (lessen) an undesired physiological condition, disorder or disease or obtain beneficial or desired clinical results. For purposes of this invention, beneficial or desired clinical results include, but are not limited to, alleviation of
30 symptoms; diminishment of extent of condition, disorder or disease; stabilized (i.e., not worsening) state of condition, disorder or disease; delay or slowing of condition, disorder, or disease progression; amelioration

-34-

of the condition, disorder or disease state, remission (whether partial or total), whether detectable or undetectable; or enhancement or improvement of condition, disorder or disease. Treatment includes eliciting a cellular response that is clinically significant, without excessive levels of side effects. Treatment also includes prolonging survival as compared to expected survival if not receiving treatment.

Conditions, diseases or disorders which may be treated with an agent of the invention include, but are not limited to, Alzheimer's disease, Huntington's disease, benign forgetfulness of memory, dementia, and other diseases associated with synaptic loss.

A pharmaceutical composition may be administered to a mammal to stimulate synaptic growth or plasticity in said mammal. A mammal refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports and pet companion animals such as a household pet and other domesticated animal such as, but not limited to, cattle, sheep, ferrets, swine, horses, poultry, rabbits, goats, dogs, cats, and the like. Preferred companion animals are dogs and cats. Preferably, the mammal is human.

A pharmaceutical composition of the invention may be administered by any suitable means, including parenteral injection (such as intraperitoneal, subcutaneous, intracranial or intramuscular injection), orally, or by topical application (e.g., transdermal or via a mucosal surface).

For oral administration, a pharmaceutical composition useful in the invention may take the form of solutions, suspensions, tablets, pills, capsules, powders, granules, semisolids, sustained release formulations, elixirs,

aerosols, and the like. Tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate are employed along with various disintegrants such as starch, preferably potato or tapioca starch, and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for generating tablets. Solid compositions of a similar type are also employed as fillers in soft and hard-filled gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the compounds of this invention may be combined with various sweetening agents, flavoring agents, coloring agents, emulsifying agents and/or suspending agents, as well as such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

The choice of formulation depends on various factors such as the mode of drug administration (e.g., for oral administration, formulations in the form of tablets, pills or capsules are preferred) and the bioavailability of the drug substance. Pharmaceutical formulations have been developed especially for drugs that show poor bioavailability based upon the principle that bioavailability may be increased by increasing the surface area, i.e., decreasing particle size. For example, U.S. Patent No. 4,107,288 describes a pharmaceutical formulation having particles in the size range from 10 to 1,000 nm in which the active material is supported on a cross-linked matrix of macromolecules. U.S. Patent No. 5,145,684

-36-

describes the production of a pharmaceutical formulation in which the drug substance is pulverized to nanoparticles (average particle size of 400 nm) in the presence of a surface modifier and then dispersed in a liquid medium to
5 give a pharmaceutical formulation that exhibits remarkably high bioavailability.

The term parenteral as used herein refers to modes of administration, which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous, intramedullary
10 and intraarticular injection and infusion. A pharmaceutical composition for parenteral injection may comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable
15 solutions or dispersions just prior to use. Aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal injection purposes. Sterile, aqueous media are readily obtainable by standard techniques well-known to those skilled in the art.
20 Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive oil), and
25 injectable organic esters such as ethyl oleate. Proper fluidity may be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

30 The pharmaceutical compositions useful in the present invention may also contain adjuvants such as, but not limited to, preservatives, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of

-37-

microorganisms can be ensured by the inclusion of various antibacterial and antifungal agents, such as for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as
5 sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents that delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the
10 drugs, it is desirable to slow the absorption from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends
15 upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming
20 microencapsule matrices of the drug in biodegradable polymers such as polylactide, polyglycolide, and polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer
25 employed, the rate of drug release may be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions that are compatible with body
30 tissues.

The injectable formulations may be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form

-38-

of sterile solid compositions which may be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Administration by slow infusion is particularly useful
5 when intrathecal or epidural routes are employed. A number of implantable or body-mountable pumps useful in delivering compound at a regulated rate are known in the art. See, e.g., U.S. Patent No. 4,619,652.

Suspensions, in addition to the active compounds, may
10 contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

For purposes of transdermal (e.g., topical)
15 administration, dilute sterile, aqueous or partially aqueous solutions (usually in about 0.1% to 5% concentration), otherwise similar to the above parenteral solutions, are prepared.

The pharmaceutical compositions useful in the
20 invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives,
25 absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

In nonpressurized powder compositions, the active ingredients in finely divided form may be used in admixture
30 with a larger-sized pharmaceutically acceptable inert carrier comprising particles having a size, for example, of up to 100 μm in diameter. Suitable inert carriers include sugars such as lactose. Desirably, at least 95% by weight

of the particles of the active ingredient have an effective particle size in the range of 0.01 to 10 μm .

The compositions useful in the present invention may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes may be used. The present compositions in liposome form may contain, in addition to the compounds of the invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art (see, e.g., Prescott (1976) *Meth. Cell Biol.* 14:33).

Other pharmaceutically acceptable carrier includes, but is not limited to, a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type, including but not limited to ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

-40-

Solid pharmaceutical excipients include, but are not limited to, starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk and the like. Liquid and semisolid excipients can be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, etc. Preferred liquid carriers, particularly for injectable solutions, include water, saline, aqueous dextrose, and glycols.

Pharmaceutical compositions useful in the present invention may contain 0.1%-95% of an agent(s) of the invention, preferably 1%-70%. In any event, the composition or formulation to be administered will contain a quantity of a agent(s) according to the invention in an amount effective to treat the condition, disorder or disease of the subject being treated.

One of ordinary skill in the art will appreciate that pharmaceutically effective amounts of the agent may be determined empirically and may be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt, ester or prodrug form. The agents may be administered to a patient as pharmaceutical compositions in combination with one or more pharmaceutically acceptable excipients. It will be understood that, when administered to, for example, a human patient, the total daily usage of the agents or composition of the present invention will be decided within the scope of sound medical judgement by the attending physician. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors: the type and degree of the cellular response

-41-

to be achieved; activity of the specific agent or composition employed; the specific agents or composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the agent; the
5 duration of the treatment; drugs used in combination or coincidental with the specific agent; and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the agents at levels
10 lower than those required to achieve the desired therapeutic effect and to gradually increase the dosages until the desired effect is achieved.

For example, satisfactory results are obtained by oral administration of agents at dosages on the order of from
15 0.05 to 500 mg/kg/day, preferably 0.1 to 100 mg/kg/day, more preferably 1 to 50 mg/kg/day, administered once or, in divided doses, 2 to 4 times per day. On administration parenterally, for example, by i.v. bolus, drip or infusion, dosages on the order of from 0.01 to 1000 mg/kg/day,
20 preferably 0.05 to 500 mg/kg/day, and more preferably 0.1 to 100 mg/kg/day, may be used. Suitable daily dosages for patients are thus on the order of from 2.5 to 500 mg p.o., preferably 5 to 250 mg p.o., more preferably 5 to 100 mg p.o., or on the order of from 0.5 to 250 mg i.v.,
25 preferably 2.5 to 125 mg i.v. and more preferably 2.5 to 50 mg i.v.

Dosaging may also be arranged in a patient specific manner to provide a predetermined concentration of the agents in the blood, as determined by techniques accepted
30 and routine in the art (HPLC is preferred). Thus patient dosaging may be adjusted to achieve regular on-going blood levels, as measured by HPLC, on the order of from 50 to 5000 ng/ml, preferably 100 to 2500 ng/ml.

The invention is described in greater detail by the following non-limiting examples.

5 Example 1: Cell Culture Preparation.

Time-mated pregnant rats were sacrificed by CO₂ asphyxiation in accordance with institutional guidelines for care and use of animals. Fetuses were removed by caesarean section and transferred to a sterile petri dish
10 with phosphate-buffered saline. Fetal hippocampi were dissected from surrounding brain tissue and meninges were completely removed. Low-density cultures of dissociated embryonic day 18 rat hippocampi (Sprague Dawley from Hilltop Laboratories, Scottsdale, PA) were prepared using
15 standard methods (Thakker-Varia et al. (2001) *supra*). Briefly, pooled tissue from each litter was mechanically dissociated in nutrient medium containing 7.5% fetal bovine serum and plated on poly-D-lysine-coated culture dishes at 350,000 cells/dish. Cultures were maintained in serum-free
20 medium (Thakker-Varia et al., 2001) for 10-14 days and contained pure neurons.

Example 2: Electrophysiological Recordings

Whole-cell patch clamp recordings were performed after 10-14 days in culture. Currents were recorded with an
25 Axoclamp 200 amplifier, digitized at 2.5 kHz with an INDEC IDA 15125 interface, filtered at 5 kHz and stored. Recording parameters and stimulus protocols were controlled by custom software written with Borland C++ that utilizes device driver libraries supplied by INDEC. Data analysis
30 programs were written with Microsoft Visual Basic. The external bath solution for voltage clamp recordings was (in mM) 1.67 CaCl₂, 1 MgCl₂, 5.36 KCl, 137 NaCl, 17 glucose, 10 HEPES and 20 sucrose (neuron recording solution, NRS). The

-43-

pipette solution contained (in mM) 105 Cs-methanesulfonate, 17.5 CsCl, 10 HEPES, 0.2 EGTA, 8 NaCl, 2 Mg-ATP, 2 Na-ATP, 0.3 Na-GTP, 20 phosphocreatinine, and 50 U/ml creatinine phosphokinase. All recordings were made at room temperature. The typical range of pipette resistance was 3-5 M Ω . Cell capacitance was 10-20 pF and access resistance was 7-20 M Ω . Vacuum perfusion system contained BDNF (50 ng/ml, Preprotech, Princeton, NJ), NRS as a control or VGF peptides (0.1 μ M = 740 ng/ml). Three synthetic rat VGF peptides were used: Leu-Glu-Gly-Ser-28 (SEQ ID NO:15) amino acids 285-312 (amidated), Thr-Leu-Gln-Pro-62 (SEQ ID NO:14) amino acids 556-617 and Ala-Gln-Glu-Glu-30 (SEQ ID NO:16) amino acids 588-617. Each data point represents a cell from a separate dish and at least three different platings were used for each condition.

Example 3: Data Analysis

Data were analyzed by integrating the synaptic currents for each sweep (synaptic charge). The charge measurements for all sweeps in a 1 min period were averaged (binned). Baseline is considered the average synaptic charge during the 2 min period (-2 to 0 min) in NRS immediately prior to BDNF or VGF application. Fold increases were then determined by dividing the synaptic charge during BDNF or VGF exposure by the baseline. Recordings were rejected if either the 0-5 min binned time period or 5-10 min binned time period after switching during BDNF or VGF exposure was 2x SEM below baseline, indicating "run down".

Example 4: Microarray

Total cellular RNA was prepared from neuronal cultures treated with either BDNF (50 ng/ml) or vehicle (water) for either 20 min or 3 hr by the guanidine isothiocyanate

-44-

method and CsCl gradients (Thakker-Varia et al., 2001). DNase-treated total RNA was prepared from neurotrophin or control hippocampal cultures and reverse transcribed using gene specific primers. Radioactively labeled probes were then hybridized to Atlas Arrays Rat 1.2 (Clontech, Palo Alto, CA) according to manufacturer's protocol. Arrays were performed in duplicate with an independent source of RNA. The hybridization pattern was analyzed on a phosphorimager and normalized to the global hybridization signal. Analysis was performed by Clontech using 1.7-fold change as criteria for significance. All transcriptional changes were confirmed by real time RT-PCR on independent samples (see below).

Example 5: Real Time RT-PCR

cDNA (100 μ l) was prepared from 2 μ g control or BDNF-treated RNA using random primers and Superscript II reverse transcriptase. 25 μ l PCR reactions were then carried out using gene specific primers designed by Primer Express software and TaqMan MGB probes (Applied Biosystems 7000 Sequence Detection System, Foster City, CA). Duplicate wells were included for each condition and primer pair. Primers specific to the housekeeping gene, *glyceraldehyde-3-phosphate dehydrogenase* (GAPDH) were used as an internal control. Data analysis was performed according to the protocol provided by Applied Biosystems.

Example 6: Single-Sell aRNA Amplification

Following whole-cell patch clamp recording for approximately 30 min, the cellular contents were aspirated into the patch pipette with a small amount of suction. cDNA synthesis was performed in vitro using the MessageAmp aRNA kit (Ambion, Austin, TX). Two rounds of aRNA synthesis were performed resulting in a one million-fold amplification of the original material (Phillips and Eberwine, 1996). 20 μ l

-45-

of cDNA was synthesized from all of the aRNA and 2 μ l was used as a template in real-time PCR for *Arc* and *GAPDH* gene expression as described above with the exception that 65 amplification cycles were used.

5 **Example 7: Pharmacological Treatments**

Hippocampal neurons were pre-incubated with inhibitors for 30 min prior to BDNF treatment (20 min for *IEGs* and 3 hr for *VGF*). All inhibitors were purchased from Calbiochem, San Diego, CA. K252a (200 nM), U0126 (30 μ M), PD98059 (50
10 μ M), U73122 (10 μ M), KN93 (10 μ M), LY294002 (30 μ M), H-89 (1 μ M) and chelerythrine chloride (CC, 1 μ M). Pharmacological inhibitors were tested for efficacy by assaying downregulation of endogenous kinase phosphorylation in cell culture using Western blot (Y.Z. Du
15 and C.F. Dreyfus, personal communication). Each experiment consisted of 4 groups: vehicle, inhibitor alone, BDNF, and BDNF plus inhibitor. Following inhibitor and BDNF treatments, RNA was isolated and subjected to real time RT-PCR as described above.

20 **Example 8: SDS-PAGE and Western Blot Analysis**

BDNF-treated and untreated hippocampal neurons plated in equal numbers were solubilized in lysis buffer (20 mM Tris pH 8, 0.5% Triton X-100, 0.5% SDS, protease inhibitor tablet, 1 mM PMSF, and 0.5 mM vanadate). The protein
25 content was determined with the BCA protein assay kit (Pierce Chemical Co, Rockford, IL). Samples containing equal amounts of protein were denatured in Laemmli's sample buffer (containing β -mercaptoethanol) for 5 min and subjected to 8- 12% gradient SDS-PAGE (Suen et al., 1997).
30 The proteins were transferred to polyvinylidene difluoride (PVDF) membranes (Millipore Corp, Bedford, MA) which were blocked for 1 hour with a 5% solution of dry milk powder plus 5% normal serum in 0.1% TWEEN 20-PBS. The PVDF

membranes were incubated with goat anti-R15 VGF (1:500) (Santa Cruz Biotechnology, Inc, Santa Cruz, CA) or goat anti-actin (1:500) (Santa Cruz) with 1% bovine serum albumin overnight at 4°C. Membranes were washed, followed by 1 h incubation with donkey anti-goat HRP-conjugated IgG (1:5000) at RT. The immunopositive bands were visualized by chemiluminescence using the ECL™ detection kit (NEN, Boston, MA), quantitated on a BIO-RAD Gel Doc and normalized to actin.

10 **Example 9: Immunocytochemistry**

Cultures were fixed in 4% paraformaldehyde. A blocking step was carried out in 30% normal horse serum in PBS containing 0.3% Triton X-100. VGF antibody (R15) (1:100) (Santa Cruz) was applied overnight at 4°C. Cultures were incubated for 1 hr at room temperature with Alexa Fluor 594 donkey anti-goat secondary (1:500) (Molecular Probe, Eugene, OR) and visualized on a Leica fluorescent microscope (DMIRB).

Example 10: Trace Eyeblick Conditioning

20 Classical eyeblink conditioning methods followed published procedures (Donahue et al., 2002; Leuner et al., 2003). Generally, an auditory stimulus is associated with stimulation to the eyelid. After repeated pairings, the animals respond to the auditory stimulus in anticipation of the eyelid stimulation. When a temporal gap is placed between the auditory stimulus and the eyelid stimulation, learning becomes hippocampal-dependent (Beylin et al., 2001). Specifically, groups of rats were exposed to 200 trials per day for 4 days for a total of 800 trials. The rats were trained with a trace paradigm in which a 250-ms 82-dB burst of white noise (CS) was separated from a 100-ms, 0.7-mA periorbital shock (US) by a 500-ms trace interval. The intertrial interval was 25 ± 5 s. During

-47-

unpaired training, rats received the same number of CS and US exposures, but presented in an explicitly unpaired manner. Naïve rats did not receive stimulus exposure. Rats were anesthetized with Nembutal (50 mg/kg) and decapitated
5 immediately after training. The hippocampus was dissected and frozen on dry ice. RNA was prepared from the tissue and subjected to real time RT-PCR as described above.

What is claimed is:

1. A method for modulating synaptic growth or plasticity comprising increasing the expression of a BDNF-inducible nucleic acid sequence or activity of a protein encoded thereby, so that synaptic growth or plasticity is stimulated.

2. The method of claim 1 wherein the nucleic acid sequence which is inducible by BDNF comprises c-fos proto-oncogene (SEQ ID NO:1); early growth response protein 1 (SEQ ID NO:2); activity-regulated cytoskeletal associated (SEQ ID NO:3); fos-related antigen 2 (SEQ ID NO:4); G1/S-specific cyclin D1 (SEQ ID NO:5); voltage-gated potassium channel protein (SEQ ID NO:6); sodium channel, beta 1 subunit (SEQ ID NO:7); secretogranin II precursor (SEQ ID NO:8); somatostatin receptor 4 (SEQ ID NO:9); transmembrane receptor UNC5 homology (SEQ ID NO:10); neuropeptide Y (SEQ ID NO:11); VGF protein precursor (SEQ ID NO:12); or protein-tyrosine phosphatase 1B (SEQ ID NO:13).

3. A method for identifying an agent which increases synaptic growth or plasticity comprising contacting a test cell with an agent and detecting activation of:



c-fos proto-oncogene (SEQ ID NO:1);
early growth response protein 1 (SEQ ID NO:2);
activity-regulated cytoskeletal associated (SEQ ID NO:3);
fos-related antigen 2 (SEQ ID NO:4);
G1/S-specific cyclin D1 (SEQ ID NO:5);
voltage-gated potassium channel protein (SEQ ID NO:6);
sodium channel, beta 1 subunit (SEQ ID NO:7);
secretogranin II precursor (SEQ ID NO:8);

-49-



somatostatin receptor 4 (SEQ ID NO:9);
transmembrane receptor UNC5 homology (SEQ ID NO:10);
neuropeptide Y (SEQ ID NO:11);
VGF protein precursor (SEQ ID NO:12); or

5 protein-tyrosine phosphatase 1B (SEQ ID NO:13)
nucleic acid sequences in the test cell wherein an increase
in the activation of said nucleic acid sequences in the
test cell contacted with the agent relative to the
activation of said nucleic acid sequences in a test cell
10 not contacted with the agent is indicative that said agent
increases synaptic growth or plasticity.

4. A method for treating a disease or condition
associated with damaged or diseased synapses comprising
15 administering an effective amount of an agent identified by
the method of claim 3.

*FIG. 1A**FIG. 1B*

2/2

*FIG. 1C**FIG. 2*

SEQUENCE LISTING

<110> University of Medicine and Dentistry of New Jersey
Black, Ira B.

<120> A METHOD FOR INCREASING SYNAPTIC GROWTH OR PLASTICITY

<130> UMD-0016

<150> US 60/422,986

<151> 2002-11-01

<160> 16

<170> PatentIn version 3.1

<210> 1

<211> 2116

<212> DNA

<213> Rattus rattus

<400> 1

cgcaactgag aagactggat agagccggcg gagccgcgaa cgagcagtga ccgcgctccc	60
accagctct gctctgcagc tcccaccagt gtctaccctt ggaccctcg ccgagctttg	120
cccaaaccac gaccatgatg ttctcgggtt tcaacgcgga ctacgaggcg tcatcctccc	180
gctgcagtag cgctccccg gccggggaca gcctttccta ctaccattcc ccagccgact	240
ccttctccag catgggctcc cctgtcaaca cacaggactt ttgcgcagat ctgtccgtct	300
ctagtgccaa ctttatcccc acggtgacag ccatctccac cagcccagac ctgcagtggc	360
tggtgcagcc cactctggte tcctccgtgg ccccatcgca gaccagagcg ccccatcctt	420
acggactccc caccctgtcg accggggctt acgccagagc gggagtgggtg aagaccatgt	480
caggcggcag agcgcagagc atcggcagaa ggggcaaagt agagcagcta tctcctgaag	540
aggaagagaa acggagaatc cgaagggaaa ggaataagat ggctgcagcc aagtgccgga	600
atcggaggag ggagctgaca gatacgtcc aagcggagac agatcaactt gaagacgaga	660
agtctgcgtt gcagaccgag attgccaatc tactgaaaga gaaggaaaaa ctggagttta	720
ttttggcagc ccaccgacct gcctgcaaga tccccaatga cctgggcttc ccagaggaga	780
tgtctgtgac ctccctggac ttgactgggg gtctgcctga ggctaccacc ccagagtctg	840
aggaggcctt caccctgcct cttctcaatg accctgagcc caagccatcc ttggagccgg	900
tcaagaacat tagcaacatg gagctgaagg ctgaaccctt tgatgacttc ttgtttccgg	960
catcatctag gccagtggc tcggagactg cccgctctgt gccagatgtg gacctgtctg	1020
gttccttcta tgcagcagac tgggagcctc tgcacagcag ttccctgggg atggggccca	1080

tggtcacaga gctggagccc ctgtgcactc ccgttggtcac ctgcactccc agctgcacta 1140
 cctatacgtc ttcctttgtc ttcacctacc ccgaggctga ctccctccct agctgcgcag 1200
 ctgcccaccg aaagggcagc agcagcaacg agccctcctc tgactcactg agctcgccca 1260
 cactgctagc cctgtgagca gtcagagaag gcagggcagc cggcactgac tgagctggtg 1320
 cattacagag aggagaaaca cgtcttccct cgaggggttc ccgtagacct agggaggacc 1380
 ttatctgtgc gtgaaacaca ccaagctgtg gacctcaagg acttgaaagc atccacatct 1440
 ggactccagt cctcacctct tccggagatg tagcaaaaaa acaaaaaaac aaaacaaaaa 1500
 aaaaacaaaa caaaaaatca aaagcaaccg catggagtgt attgtttgta gtgacacctg 1560
 agagctggta gttagtagca tgtgagccag gcctgggtct gtgtctcttt tctctttctc 1620
 cttagtcttc tcatagcatt aactaatctg ttgggttcat tattggaatt aacctggtgc 1680
 tggatatattt tcgatttgta tctagtgcag ctgattttaa caatacctac tgtgttctctg 1740
 gcaatagtgt gttccaattt agaaatgacc aatattaaac taagaaaaga tagaacttta 1800
 ttttccggta gatagaaata aatcgctata tccacgtact gtagctcttc agcgtocatg 1860
 ttcattgtca tgtaactgat catgcattgt tgaggtggtc tgaatgttct gacattaaca 1920
 gttttccatg aaaacgtttt attgtgtttt caatttatatt attaatggtg attctcagat 1980
 atttatatatt ttattttatt tttttctatc ctgaggtctt tcgacatgtg gaaagtgaat 2040
 ttgaatgaaa aaattttaag cattgtttgc ttattgttcc aagacattgt caataaaagc 2100
 atttaagttg aatgcg 2116

<210> 2
 <211> 3112
 <212> DNA
 <213> Rattus norvegicus

<400> 2
 cgcagaactt ggggagccgc cgccgcgatt cgccgccgcc gccagcttcc gccgccgcaa 60
 gatcggcccc tgccccagcc tccgcggcag ccctgcgtcc accacgggcc gcggccaccg 120
 ccagcctggg ggcccaccta cactccccgc agtgtgcccc tgcacccgc atgtaacccg 180
 gccaacatcc ggcgagtgtg ccctcagtag cttcggcccc gggctgcgcc caccacccaa 240
 catcagctct ccagctcgca cgtccgggat ggcagcggcc aaggccgaga tgcaattgat 300
 gtctccgctg cagatctctg acccgttcgg ctcccttccct cactcaccca ccatggacaa 360
 ctaccccaaa ctggaggaga tgatgctgct gagcaacggg gctccccagt tcctcggtgc 420
 tgccggaacc ccagagggca gcggcggcaa taacagcagc agcagcagca gcagcagcag 480

cgggggcggt ggtgggggcg gcagcaacag cggcagcagc gctttcaatc ctcaagggga 540
gccgagcgaa caaccctacg agcacctgac cacagagtcc tttcttgaca tcgctctgaa 600
taacgagaag gcgctggtgg agacaagtta tcccagccaa actaccgggt tgccctccat 660
cacctatact ggccgcttct ccctggagcc tgcacccaac agtggcaaca ctttgtggcc 720
tgaaccctt ttcagcctag tcagtggcct tgtgagcatg accaaccctc caacctcttc 780
atcctcagcg ccttctccag ctgcttcacg gtcttctctc gcctcccaga gccaccctc 840
gagctgtgcc gtgccgtcca acgacagcag tcccatttac tcagctgcac ccacctttcc 900
tactcccaac actgacattt ttcttgagcc ccaaagccag gcctttcctg gctctgcagg 960
cacagccttg cagtaccgcg ctctgccta cctgccacc aagggtggtt tccaggttcc 1020
catgatccct gactatctgt ttccacaaca acaggagac ctgagcctgg gcaccccaga 1080
ccagaagccc ttccagggtc tggagaaccg taccagcag ccttcgctca ctccactatc 1140
cactatcaaa gccttcgcca ctcagtcggg ctcccaggac ttaaaggctc ttaataaac 1200
ctaccagtcc caactcatca aaccagccg catgcgcaag taccccaacc ggcccagcaa 1260
gacaccccc catgaacgcc cgtatgcttg cctgttgag tcctgcgatc gccgcttttc 1320
tcgctcgat gagcttacac gccacatccg catccataca ggccagaagc ccttccagt 1380
togaatctgc atgcgtaatt tcagtcgtag tgaccacctt accaccaca tccgcacca 1440
cacaggcgag aagccttttg cctgtgacat ttgtgggaga aagtttgcca ggagtgatga 1500
acgcaaggag cataccaaaa tccacttaag acagaaggac aagaaagcag acaaaagtg 1560
cgtggcctcc tcagctgcct cttccctctc ttctaccca tcccagtggt ctacctccta 1620
cccatcccc gccaccacct catttccatc cccagtgcc acctcttact cctctccggg 1680
ctcctctacc taccgtctc ctgcacacag tggttccca tcgccctcg tggccaccac 1740
ctatgcctcc gtcccacctg cttccctgc ccaggtcagc accttccagt ctgcaggggt 1800
cagcaactcc ttcagcacct caacgggtct ttcagacatg acagcaacct tttctcctag 1860
gacaattgaa atttgctaaa gggaatgaaa gagagcaaag ggaggggagc gcgagagaca 1920
ataaaggaca ggaggggaaga aatggcccg aagagggggt gcctcttagg tcagatggaa 1980
gatctcagag ccaagtcctt ctagttagta gaaggccgt tggccaccag ccctttcact 2040
tagcgtccct gccctccca gtcccggtc ttttgacttc agctgcctga aacagccag 2100
tccaagttct tcacctctat ccaaaggact tgatttgcag ggtattggat aaaccatttc 2160
agcatcatct ccaccacatg cctggccctt gctcccttca gcactagaac atcaagttgg 2220

```

ctgaaaaaaaa aaatgggtctt gggccctcag aaccctgcc tgtatctttg tacagcatct 2280
gtgccatgga ttttgttttc cttgggggtat tcttgatgtg aagataattt gcatactcta 2340
ttgtactatt tggagttaaa ttctcacttt gggggagggg gagcaaagcc aagcaaacca 2400
atggtgatcc tctattttgt gatgatcctg ctgtgacatt aggtttgaaa cttttttttt 2460
tttttgaagc agcagtccta ggtattaact ggagcatgtg tcagagtgtt gttccgttaa 2520
ttttgtaaat actgctcgac tgtaactctc acatgtgaca aaatacgggt tgtttgggtg 2580
ggttttttgt tgtttttgaa aaaaaaattt tttttttgcc cgtccctttg gtttcaaaag 2640
tttcacgtct tgggtgccttt gtgtgacaca ccttgccgat ggctggacat gtgcaatcgt 2700
gaggggacac gctcacctct agccttaagg gggtaggagt gatgtttcag gggaggcttt 2760
agagcacgat gaggaagagg gctgagctga gctttgggtc tccagaatgt aagaagaaaa 2820
atttaaaaca aaaatctgaa ctctcaaaag tctatttttt taactgaaaa tgtagattta 2880
tccatgttcg ggagttggaa tgctgcgggt acctactgag taggcggtga cttttgtatg 2940
ctatgaacat gaagttcatt attttgtggt tttattttac ttcgtacttg tgtttgctta 3000
aacaagtga cttgtttggc ttataaacac attgaatgcg ctttactgcc catgggatat 3060
gtggtgtgta tccttcagaa aaattaaaag gaaaataaag aaactaactg gt 3112

```

```

<210> 3
<211> 3032
<212> DNA
<213> Rattus norvegicus

```

```

<400> 3
agtgtctctg cgagtagtcc tccctcagcc gcagtctctg ggcctcttca gcttgagcgg 60
cggcgagcct gccacactcg ctaagctcct ccggcaccgc gcacttgcca ctgccactgc 120
cgcttcgcgc ccgctgcagc cgccggctct gaatccttct ggcttcgcgc tcagaggagt 180
tcttagcctg tcccgaaccg taaccccggc gagcagatgg agctggacca tatgacgacc 240
ggcggcctcc acgcctaccc tgccccgcgg ggtggggcgg ccgccaaacc caatgtgatc 300
ctgcagattg gtaagtgccg agctgagatg ctggagcacg tacggaggac ccaccggcat 360
ctgttgaccg aagtgtccaa gcaggtggag cgagagctga aagggttgca caggtcgggtg 420
ggcaagctgg agaacaactt ggacggctat gtgccacgg gcgactcaca gcgctggaag 480
aagtccatca aggcctgtct ctgccgctgc caggagacca tcgccaacct ggagcgctgg 540
gtcaagcgtg agatgcacgt gtggaggagg gtcttctacc gtctggagag gtgggcccgc 600
cgcttgaggt ccatgggcgg caagtaccca gtgggcagcg agccggcccg ccacactgtc 660

```


tctgtaggtg	tggggggtcc	agagccctac	tgccaggaag	ctgatggcta	cgactacact	720
gttagccct	atgccatcac	cccgccacct	gccgcaggag	agctgcctga	gcaggagtca	780
gttggggctc	agcaatacca	gtcttggttg	ccaggtgagg	atgggcaacc	aagcccaggt	840
ctggataccc	agatctttga	ggaccacagg	gagttcctga	gccacctgga	agagtacctg	900
cggcaggtgg	gtggctctga	agaatattgg	ctgtcccaga	tccagaacca	catgaatggg	960
ccagccaaga	agtgggtgga	gttcaaacag	ggctcgggtga	agaactgggt	ggagttcaag	1020
aaggagtttc	tgcagtacag	tgagggtacg	ctctcccgcg	aagccattca	gcgggagctg	1080
gacctgccac	agaagcaggg	tgagccactt	gaccagttcc	tctggcgtaa	gcgggacctg	1140
taccagacac	tgtatgtgga	cgtgaggag	gaggagatca	ttcagtatgt	ggtgggcacc	1200
ctgcagccca	agttcaagcg	ctttctgcgc	caccacttcc	ccaagaccct	ggagcagctc	1260
atccagaggg	gcatggaagt	tcaggacggc	ctggagcagg	cagctgagcc	ttctgtcacc	1320
cctctgccca	cagaggatga	gactgaggca	ctcacgcctg	ctcttaccag	cgagtacagta	1380
gccagtgaca	ggaccagcc	tgaatagagg	ggccagccca	gggtcccag	cctgcctgcc	1440
acaccagtc	tgtggctttt	gtcaactagg	acttgattga	gctggggctg	acaccaagg	1500
ggatgccctg	tccagccaga	caccttctca	cccactggcc	tgactcaca	ctgccacaca	1560
accatgattc	atggacatca	agaagcccct	ctcccatagg	gctcccacct	gccacctacc	1620
cctcacctgt	ctgccctagt	cctggccctg	tctccagtgg	cctcacctc	tacactctca	1680
gaccatcaca	gaacaccttt	ggcttctctca	ttctgcatca	gtgtccaggg	ccctttgggt	1740
agtcaagaaa	tcaagtgtct	gaaaggcaat	gaaaagtagg	caccaaacc	aaggggcatc	1800
ccagggcaga	tgctaaagca	gaatcagaga	tggccgaagg	aacctctact	tccggggatg	1860
cagcccgtc	ctacagacac	agcagatcca	gctgggtgcc	tacctgcctc	ccagagcaac	1920
tggccagtct	tgggcagcat	agctcccctc	tcagggtgag	ctgaagcagc	agacctgacg	1980
cgtggcgcc	tcttgcccc	cagcagtgat	tcataccagt	gaagaaaagc	agacttcggc	2040
tccatgactc	agccatgcc	ggcggagggt	cccagagggg	ctgagtcctc	agccccagct	2100
gaggcagcag	ctggagtctt	cagagccagg	tgaatgacac	caggtctcaa	gctgctgaga	2160
agtctttccg	gccatgtctg	gaaggggtac	caccccagca	ccagcacctg	cccctcctct	2220
cttgaagctg	cctgcacaga	ggttccaaga	cactttcaag	gcagagaaaa	taggattaca	2280
aagaggaggt	gccttggcag	agggcagcac	ccagctcagc	ctcagagctg	aaggtgaaga	2340
caagccagcg	tgaacccccg	ggtctgccac	gaatgcccgc	tccgtggcc	actcaccagc	2400

```

tgcttgccac aagccactgc agcttgagca ggggtctgtgc cctctcagca cagagcccag 2460
ttcgctgcgt ggcctttggc ccccgccaga accttgccagg agccttaagg ttcggggcct 2520
agcccagcct gaccttacct gctgtgccct gcctgctgggt caagtccagt cccaggagac 2580
cccatgcctt ggctcctagg ctgttccagg cacttccctg acctgccggg tgattgccc 2640
gctggaacct catccacacc ccagcaccaa ccacctcgtg ttggtaactg ctctgtgtctg 2700
tagtctgagt aggccatggt gaggttcctc catctgcctg gtccattggt gttctgagac 2760
cagttocact gctgttctga cagatcccc accctgtgcc cctgccagcc cccacagggt 2820
tatttttgca cataaaccat gaccatact aatttggtta gctctgggga ctagggagac 2880
cctggagatc tcaagagtgt ggctatcccc tattttcacc aagccttcaa tatccagcca 2940
ggccatctgg cccacaccat cttacctcaa agacagacat atatatatat atacatatat 3000
atgattttgt taataaaaact atgaaattta aa 3032

```

```

<210> 4
<211> 984
<212> DNA
<213> Rattus norvegicus

```

```

<400> 4
atgtaccagg attatcccgg gaactttgac acctcgtccc ggggcagcag tggctctcct 60
gcgcacgcgc agtcctactc cagcgggtggc ggcggccagc agaagttccg ggtagatatg 120
cctggctcag gcagtgccct catccccaca atcaacgcc taccaccac cagccaggac 180
ctgcagtgga tggtagagcc cacagtgatt acctccatgt ccaacccta tccgcgctcg 240
caccctaca gcccctgcc aggtttgcga tcagtccctg ggcacatggc tctcccaga 300
ccaggagtga tcaagaccat tggtagacc gtgggcccga gaaggagaga tgagcagctg 360
tctcccgaag aggaggagaa gcgtcgaatc cggcgagaga gaaacaagct ggccgctgcc 420
aagtgtcgga accgtcgacg tgagctgacg gagaagctgc agacggagac agaggagctg 480
gaggaggaga agtcgggtct gcagaaagaa atcgctgagc tgcagaagga gaaggagaag 540
ctggagttca tgctgggtggc tcacggccct gtgtgcaaaa tcagtcctga ggaacgccga 600
tcttccccca cctccggggg acagtcctta cgcggtacgg gcagtgtgtg cgccctgtg 660
gtggtgaaac aggagcctcc tgaagaggac agcccctctt cctcagcagg gatggacaag 720
accagcgct ctgtcatcaa gccattagc attgctgggg gtggtttcta cggggaagag 780
cctctgcaca ccccatcgt ggtgacctcc acacctgcca tcaactcggg cacttcaaac 840
cttgtcttca cctaccccag tgtcctggag caggagtcgc ctgcgtcgcc ctccgagtc 900

```

tggtccaagg ctcaccgcag aagcagtagc agtggggacc agtcatcaga ctccttgaac 960
tccccactc tgctggctct gtaa 984

<210> 5
<211> 3751
<212> DNA
<213> Rattus norvegicus

<400> 5
agaaagagag ctacagactc cgcgcgctcc ggaagaccgg gtcgtagagc tcgagacaac 60
tcccgccagc agcaagacac gggagaccca accaggatga caaccctccc caggccgccc 120
gctccatgga acaccagctc ctgtgctgcg aagtggagac catccgccgc gcgtaccctg 180
acaccaatct cctcaaccga ccgggtctgc gagccatgct taagactgag gagacctgag 240
cgccctccgt ttcttacttc aagtgcgtgc agagggagat tgtgccatcc atgcggaaaa 300
tcgtggccac ctggatgcta gaggtctgcg aggagcagaa gtgcgaagag gaggtcttcc 360
cgctggccat gaactacctg gaccgtttct tgtctctgga gccctgaag aagagccgcc 420
tgcagcttct gggggccacc tgcattgtcg tggcctctaa gatgaaggag accattcccc 480
tgactgccga gaagttgtgc atctacactg acaactctat ccgccccgag gagttgctgc 540
aaatggaact gcttctggtg aacaagctca aatggaacct ggccgccatg actccccacg 600
atttcatoga acacttcctc tccaaaatgc cagaggcgga tgagaacaag cagatcatcc 660
gcaaacatgc acagaccttt gtggccctct gtgccacaga tgtgaagttc atttccaacc 720
caccctccat ggtggctgct gggagtgtgg tggccgcgat gcagggcctg aacctgggca 780
gccccaaaca cttcctctcc tgctaccgca caacgcactt tctttccaga gtcattcaagt 840
gtgacccgga ctgcctccgt gcctgccagg aacagattga agcccttctg gagtcaagcc 900
tgcgccaggc ccagcagaac atcgatccca aggccaccga ggaggaaggg gaagtggagg 960
aagaagctgg tctggcttgc acaccaccg acgtccgaga tgtggacatc tgagggccca 1020
ccgggcaggc gggagtcacc aagtagtggc atccgcgaag agaaaggagc cagcccggt 1080
gctcctgacg aggtccccct tggggatgtg ttgttaccag aaggggaagt tttgttctct 1140
ttgttggttg ttttccctta atctttctcc tttctatctg atttaagcaa aagagaaaaa 1200
aatacctgaa agctgtctta aagagagaga gagagataga attcgcatca ccctgagtat 1260
agggagacgg gggggtgcct acaaaaatag aattctgtac ccagtaatc aactagtttt 1320
ctattaatgt gcttgtctgt tctaagaata ggattaacac acaggaagtc ttgagaagga 1380
tttttgattc ttttatgtgt ttaagaaaaa aaagcttaag aaacattgct ttaaaaaaag 1440

aaggaaaaaa aatacagcaa accattgtta aagtagaagg gtttttaggta gagaaatgta 1500
ctctgctttg gtgaaaaggc acagatttgg gaaggctctc aacctcactc ccctgcttgt 1560
tcagtgccta cagacctgtt acctgatgat acctgtgctt tatcccagga gtggggcaga 1620
cctcttaacc ttactgatgg tcgattcgac ctctagcggg ctcgtagggg tggcaacctc 1680
cagggtcat gttatgggcc ctctgcccc cacaacctgc aggttcacag agcgccagcc 1740
agcacgcagg ggtagggatg aaatagtgac ataatatatt ctatttttgg ggaaccttcc 1800
tgttttgggg gttcctgttt agagagatgc tggtttttgg ctgggtggcc tgcagcccac 1860
tcaacatcag gttgaacca cagcttttct tttgtgtgtg gtttgttttg ttatgttttt 1920
ccttctccat gttcccaaaa ccattccatt tcaaagcact tttggtcagc tagctggagg 1980
cagtgttgct ggggtgtgtg gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtcggt 2040
gggggagggg gtctagtggg atggttgggg atgtccacac actcaattct gatggctgca 2100
cagcaggtgt gtagggctgg tagcatgagg tgcttgggaa gttgtgttgg gtcgagaaga 2160
gaaagctctg ttctcgtacc accgggatct tcctgcaaag tcggagggat cttttggtgc 2220
cagctggcgt ttggaagtag ggaccgtgat ggcattacct ggacaagggg attcaggacg 2280
actcttaagt ctttcacaca ggaggctttt aaacactaaa atctctaatt tatacgtaag 2340
gctacagaag agtatattt ggaagggtt cccatgacct gtgtgactca aagcaatgtg 2400
atcttccctt gattcaaacg cacaccttct tccctgctgg cgaaggtttt aggccattgt 2460
ctgagagact ggtcttttat tggccaagtg ggggtgtctc caaaaaaacc aaaccacaaa 2520
gacggagatg tgggtctctt tgactttccc aacccaattg ccccatattg agagccatcc 2580
aaacctgagg aaaattaggg gaccccaaaa gagtcttgag tctggcacat tcttgccgcc 2640
gctcccaagt tagcaacagt aggttaattg cacacctctg gctctgtgcc tttctgttca 2700
ggatgtgttg gtgggaggtg gagagcgggt ggctggagag gggatgtgag agaagaagta 2760
tggagggtag ggaccacacg ggacagaccg cggtctctt cacagcgtg ctaccaatga 2820
ctcccaggat ccccgcggtt cggaaccaga ttcacgttgc tttgtatctt tcatgttgtt 2880
ttcgtgcta ttggagggtc agttttattt tgttgttgtt ggtttgtttg ttttttttta 2940
caatgtcata aactgccatg ttcaagtttt aatttctctg tagaagagtg tatttacaga 3000
tgccctctcc gggacttttt ctatttattg tgattcaatt ttggattaat gtgattaccg 3060
ctgtattccc aacaaaaggt gttttccaca cacacacaca cacacacaca cacacacaca 3120
cacacacaca ggttctctga cacaatacct catgtatcac ctaacaatac ataggcctgc 3180

caggcgggtg gggcggtctg cctccaggga ccctgggacc ctgctgggga tcatcctgtc 3240
 atgctggggc ttcatttgat ctgggacata gcatcacagc agtcaggcca actgtgttct 3300
 gttagtattc aatattgtta cttgtagcgg cctgttgtgc atgccaccat gctgctggac 3360
 ccggagagat ttgttctgag tctctggtgc atcatataat ctgttaggtt ctagtgttct 3420
 gtcttgtttt gtgttactca cagcattgtg ctaatgtaaa gccagccgca atgctgtagg 3480
 ccccagggtc cctagcaagc tgccaaacca aaagggtcac caccagctta gctgaggcgt 3540
 cccaaccagg caggaccctt gagggctgct gtgtccatgg tgatggggtg aagttttggc 3600
 caaagggcc aaggctggtg gatccacaca gtctgccctg tgacatgaat ggctttgagg 3660
 ggctctggct ggtggtcagg ttggctttgt gtattctggt tgacacacca tggcgcttcc 3720
 cagcacagac atgtgaccag catggtccag g 3751

<210> 6

<211> 1746

<212> DNA

<213> Rattus norvegicus

<400> 6

tcctggcctc ctaccctgc accctgcac catcatgacg gtgatgtcag gggagaatgc 60
 agacgaggct tcggccgctc caggtcaccc ccaggatggc agctacccaa ggcaggcgga 120
 ccacgacgac cacgaatgct gcgagcgcgt ggtgatcaac atctccgggc tgcgcttcga 180
 gacgcagctc aagactctgg ccagttccc caacacgctg ctgggcaacc cgaagaaacg 240
 catgcgctac tttgaccctc tgaggaatga gtacttcttt gaccgcaacc ggcccagctt 300
 cgatgccatc ctttattact accagtcggg ggggcgcctg cgcaggccgg tcaacgtgcc 360
 cctggacatg ttctccgagg agattaaatt ttacgagttg ggcgaggagg ccatggagaa 420
 gttccgggaa gatgagggtc tcatcaagga agaggagcgc ccctaccgc agaaggagta 480
 ccagcgccag gtgtggctgc tctttgagta tccggagagc tcaggacctg cacgggttat 540
 tgccattgta tccgtcatgg tcatcctcat ctccatagtc atcttttgcc tggagactct 600
 ccctgagctg aaggatgaca aggacttcac gggcaccatt caccgcatcg ataaccac 660
 agtcatctac acttctaaca tcttcacaga ccctttcttc attgtggaaa ccttgtgtat 720
 catctggttc tcttttgagc tgggtggtgc cttcttcgcc tgccccagca agacagactt 780
 ctttaagaac atcatgaact tcatcgacat tgtggccatc atcccttatt tcattaccct 840
 gggcacagag atagctgagc aggaggggaa tcagaagggc gagcaggcca cttccctggc 900
 catcctcagg gtcacccgtc tggttaagggt gttcagaatc ttcaaactct cccgccactc 960

caagggcctt cagatcctgg gccagaccct caaagctagt atgagggagt tagggctgct 1020
 catctttttc ctcttcattg gcgtcatact gttttctagt gcagtgtact ttgcggaggc 1080
 ggaagaagct gagtgcact tctccagtat ccccgatgct ttctgggtggg cgggtggtgctc 1140
 catgaccact gtgggatacg gtgacatgta ccctgtgaca attggaggca agatcgtggg 1200
 ctccttgtgt gccatcgctg gtgtgctgac aattgccctg cccgtacctg tcattgtgctc 1260
 caatttcaac tatttctacc accgagaaac tgagggggaa gagcaggctc agttgctcca 1320
 tgtagttct cctaacttag cctctgacag tgacctcagc cgccgcagct cctctactat 1380
 cagcaagtct gagtacatgg agatcgaaga ggacatgaac aatagcatag ccactacag 1440
 gcaggcta atcagaactg gtaactgcac cgcaactgat caaaactgcg ttaataagag 1500
 caagctcctg accgatgttt aaaaaaagca ccaggcaagc aatcaaaagc ccccaaacia 1560
 aacccttggc gactcctgtc ccactctgta gatactttac taaaaccgta gtctttgaat 1620
 gctttattta actggcaatg cactgttgca ttgtgaattt ggggggtggg caaacctgaa 1680
 gctttcaaga tcacatttaa aaaacaaaac caaccaaaaca agcaaaaaga aaaaaaaaaac 1740
 ccaaca 1746

<210> 7
 <211> 1490
 <212> DNA
 <213> Rattus norvegicus

<400> 7
 cgcagcctgg atgcgccctg tggcgcacgc acgcagcatc ccgagcctcc cgccgcgcgc 60
 gggatgcctg ctctccgggc cccggggctt ggccccggcg gtaaccggag cgggggggag 120
 cgcccccca gcagcagctg cggcgccgc gcccgggcca gtccgcgcgc gggcccatct 180
 cctgtcgccg cgctctgga cccaccgct tgcgcggcca tggggacgct gctggtctc 240
 gtgggtggcg cgggtgctgt atcctcagcc tgggggggct gcgtggaggt ggattctgag 300
 accgaggcag tgtatgggat gaccttcaaa atcctgtgta tctcctgtaa gcgtcgtagt 360
 gagaccaccg ccgagacctt cacggagtgg acctccgcc agaagggcac agaggaattt 420
 gtcaagatcc tacgctatga gaatgaggtg ctgcagctgg aggaagatga gcgctttgag 480
 ggccgtgtgg tgtggaacgg tagtcggggc accaaggacc tgcaggacct gtccatcttc 540
 atcaccaatg tcacctaaa ccactctggc gactacgaat gtcacgtcta ccgtctctc 600
 ttctttgata attacgagca caacaccagc gtctgcaaga agatccacct ggaggtggtg 660
 gacaaggcca acagagatat ggcatccatc gtgtcagaga tcatgatgta cgtgctcatt 720

gtggtgttaa ccatatggct cgtggcggag atggtgtact gctacaagaa gattgctgct 780
gccacggaag ctgctgcaca agagaatgcc tcggaatacc tggccattac ttccgagagc 840
aaagagaact gtacaggcgt ccagggtggct gaatagcgct ggctctgggc tccgcctcaa 900
ggaagagcca gcctacgggt accctccagc cctgcagtgg ggatcagccc ctggtgggta 960
ccctcccctg gcagtgggga tcagcccatac ggtctcccca gcctcacagt tctgcagtgg 1020
agccaccagg gtgggagcgg gcagggactg atcccacctc acccaccgcc tcccacctac 1080
cctcccaccg ccatgcatga tgggtgaagc aatatggccg cccaccctg cttttgctgc 1140
ctgtttgggg gagggggcgg tgaggcgagg gggcaggccc cgccccctc tttttgctga 1200
tttgacata ggccacttcc acacgcactg ccaggccagc cgccccaccc ctgcttgatg 1260
gggtgaagag gggtcgggac agggacagta gtgggcaggg ggttctgggc ctcactctcc 1320
ctcgccttcc tccggctgga cctgggggtc ccttctctgt acacctccta gccctggccc 1380
accogccctc tctcaccagc cttcaattgt ggtctcttgg gaaggcctct tcggcctcct 1440
atctttacag aagtagtttt tgttcatgaa ataaagattc ttggactcga 1490

<210> 8
<211> 2289
<212> DNA
<213> Rattus norvegicus

<400> 8
acaatataag acagaggaaa attttaagac atgactgaat cgaaggctta ccgatttggg 60
gcagttctgc ttcttatcca cttaattttc cttgtccctg gaaccgaagc agcttccttc 120
cagcgaaacc agctgcttca gaaagaacca gacctcagat tggagaatgt ccagaagttt 180
cctagtccag aaatgatcag ggctttggag tacatagaaa agctcaggca gcaggccac 240
agagaagaaa gcagcccaga ctacaatccc taccaaggca tctctgttcc ctttcaactc 300
aaagaaaacg gagaagaaaag tcacttggca gagagctcaa gggatgtcct gagtgaagac 360
gagtggatgc ggataatact tgaggctttg aggcaggctg aaaatgagcc gccatctgcc 420
ctcaaggaga acaagcccta tgccttgaat ctggagaaga acttccctgt ggacacgcct 480
gatgactatg agactcaaca atggcctgag aggaaactca agcacatgcg gttccctctc 540
atgtatgaag agaattccag ggaaaacccc ttcaaacgca caaacgaaat agtagaagaa 600
cagtacacac cccaaagtct tgctaccctg gagtctgtgt tccaagagct tgggaaactg 660
acagggccaa gcaaccagaa gcgtgagagg gttgacgagg aacagaagct ctacacggac 720
gatgaagatg acgtgtacaa gaccaacaac attgcctatg aagatgtggg cgggggagaa 780

12

gactggagtc ctatggagga gaaaatagag actcaaacc aggaagaggt gagagacagc 840
 aaagagaaca cagaaaaaaa cgaacaaatc aatgaagaga tgaaacggtc agggcacttg 900
 gggctcccag atgaaggtaa ccggaaagag agcaaagacc agctctcaga ggacgcctcc 960
 aaggatcatca cctacttgag aaggtttagtg aatgctgtgg gcagtgggag gtcccagagt 1020
 gggcaaaacg gggacagggc agccaggcctt cttgagaggc cccttgattc tcagtctatt 1080
 tatcagctga ttgaaatctc caggaatttg cagatacccc ctgaagactt aattgagatg 1140
 ctcaaagctg gggagaaacc aaatgggttg gtggagcccg agcaggatct ggagcttgct 1200
 gttgacctag atgacatccc ggaagctgac atagaccgcc cagacatgtt tcaaagtaag 1260
 acgtctccca aggggtgggta tcccaaggca cctggctgag gtatgatgga ggccttgcca 1320
 gatggcctca gtgttgaaga cattttaaat gttttaggga tggagaatgt agcaaatcag 1380
 aagtcccat atttcccaa ccaatacagc cgagacaagg ctctgctgag gcttccttat 1440
 ggtcctggga aatctagagc caaccagatt cccaaagtag cctggatccc agacgttgaa 1500
 agcagacaag cccctatga caatctgaat gataaggacc aagaattggg agagtactta 1560
 gccaggatgc tagttaagta ccctgagctc atgaatacca accagctgaa gagagtgcc 1620
 agcccaggct cctcagaaga tgacctcaa gaagaagagc agctcgagca ggccatcaag 1680
 gagcatctgg gtcaaggaag ctcccaggaa atggagaaac tggccaaggt gagcaaaagg 1740
 atccctgcag gatccctgaa gaatgaggat accccaaata gacagtacct ggatgaagat 1800
 atgtcctga aagtgctaga gtatctcaat caagaacagg cagagcaggg aagggaacat 1860
 cttgccaaac gggccatgga aaacatgtaa acagctttaa tgccaattt cccttctttt 1920
 cccaagtga atccctccc ttctcttaa gtgtgttaat ctctatcctg ttaacactgt 1980
 aatatcttta agtgatgtac aagcagatga ctccagatag ttttggggat ctgctttact 2040
 tattctgagc tgttatgttg tgtatggatg tgtataaatg ttatgactct cagattaaaa 2100
 atatgtcctt tattcaagaa agatatctat gatagtgttg actaatgtat ccaatgggtca 2160
 tggatttgac aatgctcaca tatgatgaag agtatcctat aattatcttg gaagttttta 2220
 acatttattg aattattttg ttactgtctg tagtgttttg tggagtcttg gagcaaaatc 2280
 aataaagca 2289

<210> 9
 <211> 1865
 <212> DNA
 <213> Rattus norvegicus

13

<400> 9
gttcagcggt cggctgctct ccacggcaat ccgctgcccc gggtagggcac cccgaagcat 60
gaacacgcct gcaactctgc ccctgggggg cgaggacacc acctggaccc ctgggatcaa 120
cgccagctgg gctccggatg aggaggagga tgcagtgcgg tccgacggca cggggacagc 180
gggcatggta actatccagt gcatctatgc gctcgtgtgt ctggtagggcc tggtaggaaa 240
cgccctggtc atattcgtga tcctacgcta tgccaaaatg aagacagcca ccaacatcta 300
cctgctcaac ctggccgtcg ctgatgagct cttcatgctc agtgtgccat ttgtggcctc 360
ggcggctgcc ctgcgccact ggccgttcgg ggcgggtgctg tgccgcgcag tgcttagtgt 420
ggacggcctt aacatgttca cgagtgtctt ctgcctcaca gtgctcagcg tggatcgcta 480
tgtggctgta gtgcaccctc tgcgagctgc cacctaccgg cggcccagcg tggccaagct 540
aatcaacctg ggagtgtggc tagcatcctt gctggtcacc ctgcccacgc cagtcttcgc 600
tgacactagg ccagctcgtg ggggtgaggc agtagcttgc aacctgcact ggcctcacc 660
ggcctgggtc gcagtctttg tgatctatac ttttttgctg ggcttcctac tcccggttct 720
ggctatcgga ttatgttacc tgcttatcgt gggcaagatg cgtgctgtgg cctgcggggc 780
tggttgcaa caacggaggc gctcagagaa gaagatcact aggctcgtgc taatggtggt 840
gactgtcttt gtgctatgct ggatgccatt ctatgtagtg cagcttctga atctgtttgt 900
caccagcctc gatgccactg tcaacctagt gtccctcacc ctgagctatg ccaacagctg 960
tgccaaccgc attctctatg gtttcctctc agacaacttc cgacgctctt tccagcgggt 1020
tctgtgcctg cgctgctgtc tcctggaaac aactggagggt gctgaggaag agcccctgga 1080
ctactatgct actgctctca aaagcagagg tggcccagga tgcatatgcc ctccattgcc 1140
ctgccagcag gagcccatgc aagcagaacc tgccctgcaag cgagtccctt tcaccaagac 1200
cactactttc tgaaaaccat ttcaccctcc ctgagccac ctgcaagcag gtctgcacca 1260
cactctcaag ccagcaactt caagaaaact cctgttgtca ctaagccagg ccctttcagc 1320
agcctgtgtt ctgtccctag gagcctcagg actcctgcta gccctgcct ctcttaggac 1380
tgactggctc caaggacaac tccgtggggg taggacttct ctgggttttg ggctagagta 1440
ccatccatcc tttcctggac ctctagcaat ttttcaagag gcaggaagca ggtgggtggtc 1500
agaaagggat gcctaccctt gtgtgacttg tgacagtgc tgcttggaag agcgtggga 1560
gggtgaggta ggcagagcta ggctctctgc tgttggttag catagggcat acggtgatac 1620
aggggagaag atatgatacc tccaagtgtt ttccctctgt gtctgtctga gtctctgtt 1680
gctaaatgag atgtctacgc aacagctgaa agcatttgct ttccaaggc aaatgtttct 1740

14

ccagttgtca aaggaccagt agcagacttc ctgcgaatgc aaatgtttta agaaggatgg 1800
tgtggggcgt tttttgaaaa aaaaaataat tctgatttct ggtcaggaat taaaaggcag 1860
aaagg 1865

<210> 10
<211> 2838
<212> DNA
<213> Rattus norvegicus

<220>
<221> misc_feature
<222> (2265)..(2265)
<223> n=a, c, g or t

<400> 10
atgagggccc ggagcggggc gcggggcgcg ctgctgctgg cgctgctgct ctgctgggat 60
ccgacaccga gcttagcagg cattgactct ggtggccagg cactcccaga ctccctccca 120
tcagcaccgc cggagcagct gcctcacttc ctgctggaac cagaggatgc ctacatcgta 180
aagaacaagc cagtggaatt gcactgccga gccttccttg ccacacagat ctacttcaag 240
tgtaatggcg agtgggtag ccagaaaggc cacgtcacgc aggagagcct ggatgaggcc 300
acaggcttgc gaatacgaga ggtgcagata gaggtgtcgc ggcagcaggt ggaggaactt 360
tttgggctcg aggactactg gtgtcagtgc gtggcctgga gctcttcggg aaccaccaag 420
agtcgccgag cctacatccg cattgcctac ttgcgcaaga actttgacca ggagcctctg 480
gcgaaggagg tacccttgga tcatgaggtc cttctgcagt gccgccacc agagggagtg 540
cctgtggctg aggtggaatg gctcaagaat gaagatgtca tcgatccgc tcaggacact 600
aacttcctgc tcaccattga ccacaacctc atcatccgcc aggcgcgcct ctccagacaca 660
gccaaactaca cctgtgtggc aaagaatatt gtggccaagc gccggagcac gacggccaca 720
gtcatcgtct atgtgaacgg aggttggtcc agctgggcag aatgggcacc ctgctctaac 780
cgctgcggcc gaggttgga gaaacgtact aggacctgca ccaaccagc cccactcaat 840
ggaggtgcct tctgcgagg acaggcttgc cagaagacgg cttgcaccac cgtgtgcca 900
gtggatggag cgtggactga gtggagcaag tggccgcct gcagcacaga gtgtgcgcac 960
tggcgcagcc gcgagtgcac ggcaccgccg cccagaacg gaggcctga ctgcagcggg 1020
acgctacttg actccaagaa ctgcaccgat gggctgtgcy tgctgaatca gagaactcta 1080
aacgacccta aaagccgccc cctggagccg tcgggagacg tggcgctgta tgcgggcctc 1140
gtgggtggccg tctttgtggg tctggcagtt ctcattggctg taggagtgat cgtgtaccgg 1200

agaaactgcc gggacttcga cacggacatc actgactcct ctgctgccct cactggtggt 1260
ttccaccccg tcaacttcaa gactgcaagg cccagcaacc cacagctcct gcacccatcc 1320
gcccctccgg acctaacggc cagtgtctggc atctaccgcg gacctgtgta tgccctgcag 1380
gactctgccg acaagatccc tatgactaat tcaccccttc tggatccctt gccagcctc 1440
aagatcaagg tctatgactc cagcaccatc ggctctgggg ctggcctggc tgatggagcc 1500
gacctgtctg gtgtcttacc acccggtaca taccagggcg atttctcccg ggacaccac 1560
ttcctgcacc tgcgcagcgc cagccttggt tccagcacc tcctgggcct ccctcgagac 1620
cccagcagca gtgtcagtgg cacctttggt tgccctgggtg ggaggctgac cattcccggc 1680
acaggggtca gcctgttggt accaaatgga gccattcccc agggcaagtt ctatgacttg 1740
tatctacgta tcaacaagac tgaaagcacc ctcccacttt cggaagggtc ccagacagta 1800
ttgagccct cgggtgacctg cgggcccacg ggccctctcc tgtgccgcc tgttgtcctc 1860
actgtgcccc actgtgtctga agtcattgcc ggagactgga tcttccagct caagaccag 1920
gcccacaggg gccactggga ggagggtgtg actttggatg aggagactct gaacaccccc 1980
tgctactgcc agctagaggc taaatcctgc cacatcctgt tggaccagct gggtacctac 2040
gtgttcacgg gcgagtccta ctcccgtcc gcagtcaagc ggctccagct agccatcttc 2100
gccccagccc tctgcacctc cctggagtat agtctcaggg tctactgtct ggaggacact 2160
cctgcagcac tgaaggaggt cctagagctg gagaggactc tgggtggcta cttggtggag 2220
gagcccaaga ctttgctctt taaggacagt taccacaacc tacgnctctc cctccatgac 2280
atcccccacg cccactggag gagcaaacta ctggccaagt accaggagat tcccttctac 2340
catgtgtgga acggcagcca gaaagccctg cactgcactt tcaccctgga gagacatagc 2400
ctagcctcca ctgagttcac ctgtaaggtc tgcgtgcggc aggtagaagg ggaaggccag 2460
attttccagc tgcacaccac gctggctgag acgcctgctg gctccctgga tgcactctgc 2520
tctgcccctg gcaatgctgc caccacacag ctgggaccct atgccttcaa gataccactg 2580
tccatccgcc agaagatctg caacagcctg gacgccccca actcacgggg caatgactgg 2640
cggctgttgg cacagaagct ctccatggac cggtagctga actacttcgc caccaaagct 2700
agtccacag gcgtgatctt agacctctgg gaagctcggc agcaggatga tggggacctc 2760
aacagcctgg ccagtgcctt ggaggagatg ggcaagagt agatgctggt agccatgacc 2820
actgatggcg attgctga 2838

16

<210> 11
<211> 511
<212> DNA
<213> Rattus norvegicus

<400> 11
gcgcccagag cagagcaccc gctgcgcaga gaccacagcc cgcccgccat gatgctaggt 60
aaciaacgaa tggggctgtg tggactgacc ctgctctat ccctgctcgt gtgtttgggc 120
attctggctg aggggtaccc ctccaagccg gacaatccg gcgaggacgc gccagcagag 180
gacatggcca gatactactc cgctctgcga cactacatca atctcatcac cagacagaga 240
tatggcaaga gatccagccc tgagacactg atttcagatc tcttaatgag agaaagcaca 300
gaaaatgccc ccagaacaag gcttgaagac ccttccatgt ggtgatggga aatgaaactt 360
gctctcctga cttttcctag tttcccccca catctcatct catcctgtga aaccagtctg 420
cctgtcccac caatgcatgc caccaccagg ctggattccg acccatttcc cttgttgcg 480
ttgtatatat gtgtgtttaa ataaagtatc a 511

<210> 12
<211> 2556
<212> DNA
<213> Rattus norvegicus

<400> 12
ctttcatact ccagccacgg aacggagcca gggcagacgg gtccggattt tccccctgcc 60
ccgaccctcc totccacctc ccgccgtcgt gacaccggct gggggcgaca ggaggcactg 120
ggtacccaga acgaggattg cgagcgttct ctgaccattt gcacgacccc agagattgcc 180
acatctttct tgttctctgc taaacgtttc tcttcgggtct ctggcagccc gttggtcatg 240
aaaaccttca cgttgccagc atccgtcctc ttctgcttcc ttctactcat ccggggggtg 300
ggagcagcac cccccgggcg ctccgatgtt tatcctcctc ccctcggctc tgagcataat 360
gggcaggtag ctgaggacgc agtgtcccgg ccaaaggatg acagcgtccc agaggtccga 420
gcggtctgga attccgagcc tcaggaccag ggagagctct tcaggggcgt ggatccccgg 480
gcgctggccg cggtagtgtt gcaggcactg gaccgtccgg cctcgcccc ggctgtcccg 540
gcaggttccc agcaggaac acccgaagaa gcagcagaag ctctgctgac cgagtccgtg 600
cgcagtcaga cccatagcct cccggcatca gaaatccaag cgtccgctgt ggcgccccct 660
cgccctcaga ctcaggacaa cgatcccag gcagacgacc gctcagaaga gctggaggca 720
ctagcatcct tgctccaaga acttcgagat ttcagtccga gtaatgctaa gcgccagcaa 780
gagacggcgg cagcagagac tgaaaccgc acgcacacgc tgaccgagt caatctggag 840

17

agccccgggc cagagcgcggt atggcgcgct tcctgggggag agttccaggc gcgcggtcccg 900
gagcgtgctc ctctgccgcc ctcggtccct tctcaattcc aggctcgaat gtccgaaaac 960
gttccccttc ccgaaaccca tcagttcggg gaaggagtgt cctcccctaa aacacatctt 1020
ggtgagactt tgacaccctt atccaaggcg taccaaagtc taagtgcctt cttcccctaa 1080
gtgcgtcggc tcgagggctc attcctgggc ggttccgagg caggagagcg cctgcttcaa 1140
caagggttag ctccaggtaga ggcagggagg aggcaggcgg aggccacccg gcaggccgca 1200
gcgcaagaag agcggctggc cgatctcgcc tccgacctgc tgctccagta tttgctgcag 1260
ggcggcgccc ggcagcgca tctcgggggt cgcgggctgc aggagacgca gcaagagcgg 1320
gagaacgaga gggaggagga ggcggagcag gagagacgcg gtggtgggga ggacgaggtg 1380
ggggaagagg atgaggaggc ggcagaggcg gaggcggagg cagaggaggc ggagaggcgg 1440
cggcagaacg cgctcctgtt cgccgaggag gaggacgggg aagccggagc cgaggacaag 1500
cgctcccagg aggaggcgcc aggccatcgg cggaaggatg ctgaggggac agaggagggc 1560
ggggaggagg atgacgacga cgaagagatg gatccgcaga cgatcgatag tctcattgaa 1620
ctgtccacca aactccacct gccagcagac gatgtgggtc gcatcatcga agaggtggag 1680
gagaaacgga agcggaagaa gaacggccct cccgagccgg tgcgcccccc cagggtgcc 1740
ccagccccga cccatgtccg ctccccgcag cccccacctc ccgccccggc ccgggatgag 1800
ttgccggact ggaacgaagt actcccaccc tgggatcggg aggaggatga ggtgtttccc 1860
ccggggccct atcaccctt cccaaactac attcgccgc ggacactgca gccgcccgca 1920
tcctcccgcc gccgtcactt ccatcacgcg ttgccacctg cgcgccacca tcccgatctg 1980
gaggcccagg ccaggcgcg gcaggaggaa gcggacgcgg aggagcgccg gctgcaggag 2040
caggaggagc tggagaatta cattgagcac gtgctgctgc accgcccgtg accgcccct 2100
gcgcgcccgc tcccaactgc gcgcgcccgc acgcccccc tcctgtctgc tctcctccc 2160
tctcgggtgt tgcatgcgc ccggctccgc ccctcggtg ccgcccggcc ccgcccaca 2220
aggccccgcc ccgggttctg tcaggaccag acctgtcaga cttctttggg gtctgatcct 2280
ggggccagcc caggcgggtg tgtggtttgt gcgagtcccc ttacaccccc acttctcca 2340
ggggcctcgt ccccatctag tttctctagc gacttcctgg tcccaaacgg ggaaaagctg 2400
ttctatthaa tcgtgtgaag tgtctgtctc ccagccttgg ggcccccgga gcctcccttc 2460
tccaaattgc tgtgaactta ccacatctt gcccttctgt tgtaaatacc cctcacggag 2520
gaaatagttt tgctaagaaa taaaagtgac tatttt 2556

<210> 13
<211> 4127
<212> DNA
<213> *Rattus norvegicus*

<400> 13
agccgctgct ggggaggttg gggctgaggt ggtggcgggc gacgggcctc gagacgcgga 60
gcgacgcggc ctagcgcggc ggacggccga ggaactcgg gcagtcgtcc cgtcccgcga 120
tggaatgga gaaggaattc gagcagatcg ataaggctgg gaactgggcg gctatttacc 180
aggatattcg acatgaagcc agtgacttcc catgcagaat agcgaaactt cctaagaaca 240
aaaaccggaa caggtaccga gatgtcagcc cttttgacca cagtoggatt aaattgcatc 300
aggaagataa tgactatata aatgccagct tgataaaaat ggaggaagcc cagaggagct 360
atatcctcac ccagggccct ttaccaaaca cgtgcgggca cttctgggag atgggtgtggg 420
agcagaagag caggggcgtg gtcattgctca accgcatcat ggagaaaggc tcgttaaaat 480
gtgccagta ttggccacag aaagaagaaa aagagatggt cttcgatgac accaatttga 540
agctgacact gatctctgaa gatgtcaagt catattacac agtacggcag ttggagttagg 600
agaacctggc taccagagg gctcgagaga tcttgcattt cactacacc acctggcctg 660
actttggagt ccctgagtca cctgcctctt tctcaattt cctattcaaa gtccgagagt 720
caggctcact cagcccagag cacggcccca ttgtggtcca ctgcagtgtt ggcattggca 780
ggtcagggac cttctgcctg gctgacacct gcctcttact gatggacaag aggaaagacc 840
cgtcctctgt ggacatcaag aaagtgtgtt tggagatgag cagggtccgc atggggctca 900
tccagacggc cgaccaactg cgcttctcct acctggctgt gatcgagggt gcaaagttca 960
tcatgggcca ctctgcagtg caggatcagt ggaaggagct tcccatgaa gacctggagc 1020
ctccccctga gcacgtgccc ccacctcccc ggccacccaa acgcacattg gagcctcaca 1080
atggcaagtg caaggagctc ttctccaacc accagtgggt gagcgaggag agctgtgagg 1140
atgaggacat cctggccaga gaggaagca gagccccctc aattgctgtg cacagcatga 1200
gcagtatgag tcaagacact gaagttagga aacggatggt ggggtggaggt cttcaaagtg 1260
ctcaggcatc tgtccccact gaggaagagc tgtcccaac cgaggaggaa caaaaggcac 1320
acaggccagt tcaactggaag cccttctctg tcaactgtgt catggccacg gccctggcga 1380
ctggcgcgta cctctgttac cgggtatgtt ttactgaca gactgctgtg aggcattgagc 1440
gtgggtggcg ctgccactgc ccaggttagg atttggctct cggcgtctaa cctgggtgtg 1500
aagaaacaac agcttacaag cctgtggttg aactggaagg gccagcccca ggaggggcat 1560

ctgtgcactg ggctttgaag gagcccctgg tccaagaac agagtctaata ctcagggcct 1620
taacctgttc aggagaagta gaggaaatgc caaatactct tcttgctctc acctcactcc 1680
tcccctttct ctggttcgtt tgttttttggg aaaaaaaaaa aaagaattac aacacattgt 1740
tgtttttaac atttataaag gcaggttttt gttattttta gagaaaacaa aagatgctag 1800
gcactggtga gattctcttg tgccctttgg catgtgatca gattcacgat ttacgtttat 1860
ttccggggga ggggtcccacc tgtcaggact gtaaagttcc tgotggcttg gtcagccccc 1920
ccaccccccc accccgagct tgcaggtgcc ctgctgtgag gagagcagca gcagaggctg 1980
cccctggaca gaagcccagc tctgcttccc tcaggtgtcc ctgcgtttcc atcctccttc 2040
tttgtgaccg ccatcttgca gatgaccag tcctcagcac cccaccctg cagatgggtt 2100
tctccgaggg cctgcctcag ggtcatcaga ggttggtgc cagcttagag ctggggcttc 2160
catttgattg gaaagtcatt actattctat gtagaagcca ctccactgag gtgtaaagca 2220
agactcataa aggaggagcc ttggtgtcat ggaagtcact ccgcgcgcag gacctgtaac 2280
aacctctgaa acactcagtc ctgctgcagt gacgtccttg aaggcatcag acagatgatt 2340
tgcagactgc caagacttgt cctgagccgt gatttttaga gtctggactc atgaaacacc 2400
gccgagcgt tactgtgcag cctctgatgc tggttggctg aggtgcggg gaggtggaca 2460
ctgtgggtgc atccagtga gttgcttttg tgcagtggg tccagcagca cagcccgcac 2520
tccagcctca gctgcaggcc acagtggcca tggaggccgc cagagcgagc tgggggtgat 2580
gcttgctcac ttggagcagc cttcccagga cgtgcagctc ccttctgtct ttgtccttct 2640
gcttccttcc ctggagtagc aagcccacga gcaatcgtga ggggtgtgag ggagctgcag 2700
aggcatcaga gtggcctgca gcggcgtgag gcccttccc ctccgacacc cccctccaga 2760
ggagccgctc cactgttatt tattcacttt gccacagac acccctgagt gagcacacc 2820
tgaaactgac cgtgtaagggt gtcagcctgc acccaggacc gtcaggtgca gcaccgggtc 2880
agtcctaggg ttgaggtagg actgacacag ccaactgtgtg gctggtgctg gggcaggggc 2940
aggagctgag ggtcttagaa gcaatcttca ggaacagaca acagtgggtga catgtaaagt 3000
ccctgtggct actgatgaca tgtgtaggat gaaggctggc ctttctccca tgactttcta 3060
gatcccgctc cccgtctgtt ttccctgtga gtagaaaaac acacaggctc ctgtcctggt 3120
ggtgccgtgt gcttgacatg ggaaacttag atgcctgtct actggcgggc acctcgcat 3180
cgccaccact cagagtgaga gcagtgtgt ccagtgccga ggccgcctga ctcccgagc 3240
gactcttcag gctctggcct gcccagcac acccgcgtgg atctcagaca ttccacacc 3300

20

acacctcatt ccctggacac ttgggcaagc aggcccgccc ttccacctct ggggtcagcc 3360
cctccattcc gagttcacac tgctctggag caggccagga ccggaagcaa ggcagctggt 3420
gaggagcacc ctctgggaa cagtgtaggt gacagtcctg agagtcagct tgctagcgct 3480
gctggcacca gtcaccttgc tcagaagtgt gtggctcttg aggctgaaga gactgatgat 3540
ggtgctcatg actcttctgt gaggggaact tgaccttcac attgggtggc tttttttaa 3600
ataagcgaag gcagctggaa ctccagtctg cctcttgcca gcacttcaca ttttgccttt 3660
caccagaga agccagcaca gagccactgg ggaaggcgat ggccttgcct gcacaggctg 3720
aggagatggc tcagccggcg tccaggctgt gtctggagca gggggtgcac agcagcctca 3780
cagggtgggg cctcagagca ggcgctgcc tgtccctgc ccgctggag gcagcaaagc 3840
tgctgcatgc cttaagtcaa tacttactca gcagggcgct ctogttctct ctctctctct 3900
ctctctctct ctctctctct ctctctctct ctctaaatgg ccatagaata aaccatttta 3960
caaaaataaa agccaacaac aaagtgtctt ggaatagcac ctttgcagga gcgggggggtg 4020
tctcagggtc ttctgtgacc tcaccgaact gtccgactgc accgtttoca acttgtgtct 4080
cactaatggg tctgcattag ttgcaacaat aaatgttttt aaagaac 4127

<210> 14
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 14

Thr Leu Gln Pro
1

<210> 15
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 15

Leu Glu Gly Ser
1

21

<210> 16
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 16

Ala Gln Glu Glu